

PATENT APPLICATION

CANNABINOID RECEPTOR LIGANDS

INVENTORS:

Joseph A. Kozlowski, a citizen of the United States of America, residing at 1066 Stuart Road Princeton, New Jersey 08540, U.S.A.

Neng-Yang Shih, a citizen of the United States of America, residing at 1 Maple Drive, North Caldwell, New Jersey 07006, U.S.A.

Brian J. Lavey, a citizen of the United States of America, residing at 25 Hickory Place, Apt. H-10, Chatham, New Jersey 07928, U.S.A.

Razia K. Rizvi, a citizen of the United States of America, residing at 22 Tomar Court, Bloomfield, New Jersey 07003, U.S.A.

Bandarpalle B. Shankar, a citizen of the United States of America, residing at 1124 Van Arsdale Drive, Branchburg, New Jersey 08853, U.S.A.

James M. Spitler, a citizen of the United States of America, residing at 316 Wells Street, Westfield, New Jersey 07090, U.S.A.

Ling Tong, a citizen of China, residing at 8 Hemlock Circle, Warren, New Jersey 07059, U.S.A.

Ronald L. Wolin, a citizen of the United States of America, residing at 16309 Los Rosales Street, San Diego, California 92127, U.S.A.

Michael K. Wong, a citizen of Canada, residing at 999 Hidden Lake Drive, Apt. 3D, North Brunswick, New Jersey 08902, U.S.A.

ASSIGNEE: Schering Corporation

"Express Mail" Label No. EV 024688227 US

Date of Deposit: February 6, 2002

Elliott Korsen
Schering-Plough Corporation
Patent Department, K-6-1, 1990
2000 Galloping Hill Road
Kenilworth, New Jersey 07033-0530
Telephone No.: (908) 298-5684
Facsimile No.: (908) 298-5388

CANNABINOID RECEPTOR LIGANDS

BACKGROUND OF THE INVENTION

This invention relates to cannabinoid receptor ligands and, more particularly, to compounds that bind to cannabinoid (CB₂) receptors. Compounds according to the present invention generally exhibit anti-inflammatory and immunomodulatory activity and are useful in treating conditions characterized by inflammation and immunomodulatory irregularities. Examples of conditions which may be treated

5 include, but are not limited to, rheumatoid arthritis, asthma, allergy, psoriasis, Crohn's disease, systemic lupus erythematosus, multiple sclerosis, diabetes, cancer, glaucoma, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, and nephritis. The invention also relates to pharmaceutical compositions containing said compounds.

10 Cannabinoid receptors belong to the superfamily of G-protein coupled receptors. They are classified into the predominantly neuronal CB1 receptors and the predominantly peripheral CB2 receptors. While the effects of CB1 receptors are principally associated with the central nervous system, CB2 receptors are believed to have peripheral effects related to bronchial constriction, immunomodulation and

15 inflammation. As such, a selective CB2 receptor binding agent is expected to have therapeutic utility in the control of diseases associated with inflammation, immunomodulation and bronchial constriction such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, diabetes, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, nephritis, inflammatory disorders of the lungs and


20 gastrointestinal tract, and respiratory tract disorders such as reversible airway obstruction, chronic asthma and bronchitis (see, e.g., R.G. Pertwee, Curr. Med. Chem. 6(8), (1999), 635).

Various compounds have reportedly been developed which interact with CB₂ receptors and/or which have, *inter alia*, anti-inflammatory activity associated with

25 cannabinoid receptors. See, e.g., U.S. Pat. Nos. 5,338,753, 5,462,960, 5,532,237, 5,925,768, 5,948,777, 5,990,170, 6,013,648 and 6,017,919.

SUMMARY OF THE INVENTION

This invention relates to compounds of formula I:



- 5 a prodrug thereof, or a pharmaceutically acceptable salt, solvate or stereoisomer of the compound or of said prodrug; wherein:

R¹ is H, alkyl, haloC₁-C₆ alkyl, cycloalkyl, cycloalkylNH-, arylalkyl, heterocycloalkyl, heteroaryl, N(R²)₂, or NR²aryl, unsubstituted aryl or aryl substituted

- 10 with one to three X;

R² is the same or different in each occurrence and is independently selected from H or C₁-C₆ alkyl;

R³ is H, C₁-C₆ alkyl, Cl, F, CF₃, OCF₂H, OCF₃, OH or C₁-C₆ alkoxy;


R⁴ is H, C₁-C₆ alkyl, C₁-C₆ alkoxy, cycloalkyl, alkenyl, aryl, benzyl, heteroaryl,


- 15 heterocycloalkyl, arylNH-, heteroarylNH-, cycloalkylNH-, N(R²)₂, or NR²aryl, said alkyl, alkoxy, cycloalkyl, alkenyl, phenyl or heteroaryl optionally substituted with one to three X;

R⁵ is H or C₁-C₆ alkyl;


R⁶ is H or C₁-C₆ alkyl; or


- 20 R⁵ and R⁶ taken together with the carbon atom form a carbonyl group;

L¹ is C₁-C₆ alkylene, C₂-C₆alkenylene, C(R²)₂, , -CHOR²-, NOR⁵-, -SO₂-, -SO-, -S-, -O-, -NR²-, -C(O)NR²-, -NR²C(O)-, -CHCF₃- or -CF₂-;

L² is a covalent bond, C₁-C₆ alkylene, -C(R²)₂, , -CHOR²-, -C(R²)OH, NOR⁵-, -SO₂-, -NR²SO₂-, -SO-, -S-, -O-, -SO₂NR²-, -N(R²)-, -C(O)NR² or -NR²C(O)-;

- X is the same or different, and is independently selected from H, halogen, CF₃, CN, OCF₂H, OCF₂CF₃, OCF₃, OR², C₁-C₆ alkyl, cycloalkyl, cycloalkoxy, C₁-C₆ alkoxy, 5 alkoxyC₁-C₆ alkoxy, O-cycloalkyl, cycloalkylamino, cycloalkylalkoxy, heteroalkyl, -OSO₂R², -COOR², -CON(R²)₂, NHR², arylNH-, N(R²)₂, or NR² aryl;

Y is a covalent bond, -CH₂-, -SO₂-, or ;

Z is a covalent bond, -CH₂-, -SO₂- or ; or

- Y, R¹, Z and R² can be taken together with the nitrogen atom to form a 10 heterocycloalkyl; with the proviso that if Y is a covalent bond, R¹ cannot form a N-N bond with the nitrogen atom; and n is an integer of 0 to 4.

- Cannabinoid receptor ligands according to the present invention have anti- 15 inflammatory activity and/or immunomodulatory activity and are useful in the treatment of various medical conditions including, e.g., cutaneous T cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, osteoporosis, renal ischemia, myocardial infarction, cerebral stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing aveolitis, 20 psoriasis, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive pulmonary disease (COPD) or bronchitis. It is contemplated that a compound of this invention may be useful in treating more than one of the diseases listed.

- 25 Additionally, a compound of the present invention may be co-administered or used in combination with disease-modifying antirheumatic drugs (DMARDS) such as methotrexate, azathioprine leflunomide, penicillamine, gold salts, mycophenolate mofetil, cyclophosphamide and other similar drugs. They may also be co-administered with or used in combination with NSAIDS such as piroxicam, naproxen,

indomethacin, ibuprofen and the like; COX-2 selective inhibitors such as Vioxx® and Celebrex®; COX-1 inhibitors such as Feldene; immunosuppressives such as steroids, cyclosporine, Tacrolimus, rapamycin, muromonab-CD3 (OKT3), Basiliximab and the like; biological response modifiers (BRMs) such as Enbrel, Remicade, IL-1

- 5 antagonists, anti-CD40, anti-CD28, IL-10, anti-adhesion molecules and the like; and other anti-inflammatory agents such as p38 kinase inhibitors, PDE4 inhibitors, TACE inhibitors, chemokine receptor antagonists, Thalidomide and other small molecule inhibitors of pro-inflammatory cytokine production. They may also be co-administered with or used in combination with H1 antagonists such as Claritin, Clarinex, Zyrtec,
- 10 Allegra, Benadryl, and other H1 antagonists. Other drugs that the compounds of the invention may be co-administered or used in combination with include Anaprox, Arava, Arthrotec, Azulfidine, Aspirin, Cataflam, Celestone Soluspan, Clinoril, Cortone Acetate, Cuprimine, Daypro, Decadron, Depen, Depo-Medrol, Disalcid, Dolobid, Naprosyn, Gengraf, Hydrocortone, Imuran, Indocin, Lodine, Motrin, Myochrysine,
- 15 Nalfon, Naprelan, Neoral, Orudis, Oruvail, Pediapred, Plaquenil, Prelone, Relafen, Solu-Medrol, Tolectin, Trilisate and Volataren. These include any formulations of the above-named drugs.

For the treatment of multiple sclerosis, the compounds of the invention may be co-administered or used in combination with Avonex, Betaseron, Rebif and Copaxone.

- 20 These include any formulations of the above-named drugs.

For the treatment of psoriasis, the compounds of the invention may be co-administered or used in combination with steroids, methotrexate, cyclosporin, Xanelin, Amivere, Vitamin D analogs, topical retinoids, anti-TNF- α compounds and other drugs indicated for this condition. These include any formulations of the above-named

- 25 drugs.

For the treatment of asthma, the compounds of the invention may be co-administered or used in combination with Singulair, Accolate, Albuterol, and other drugs indicated for this disease. These include any formulations of the above-named drugs.

- 30 For the treatment of inflammatory bowel disease or Crohn's disease, the compounds of the invention may be co-administered or used in combination with sulfasalazine, budesonide, mesalamine and other drugs indicated for these diseases. These include any formulations of the above-named drugs.

In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I in a pharmaceutically acceptable carrier.

DETAILED DESCRIPTION

5 Unless otherwise defined, the following definitions shall apply throughout the specification and claims.

When any variable (e.g., R²) occurs more than one time in any constituent, its definition in each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

10 "Alkyl" means straight or branched alkyl chains of 1 to 12 carbon atoms. The term includes the isomers thereof such as isopropyl, isobutyl, sec-butyl, etc.

"Haloalkyl" means alkyl having 1 or more halo atoms.

15 "Heteroalkyl" means straight or branched alkyl chain comprised of from 1 to 12 carbon atoms and 1 or more heteroatoms independently selected from the group consisting of N, O and S.

"Cycloalkyl" as used herein refers to an aliphatic ring system having 3 to 10 carbon atoms and 1 to 3 rings, including, but not limited to cyclopropyl, cyclopentyl, cyclohexyl, norbornyl, and adamantyl among others.

20 "Heterocycloalkyl" means cycloalkyl containing one or more heteroatoms.

"Aryl" means an aromatic monocyclic or polycyclic ring system comprising from 6 to 14 carbon atoms. Non-limiting examples include phenyl, naphthyl, indenyl, tetrahydronaphthyl and indanyl.

25 "Arylalkyl" means a group in which the aryl and alkyl are as previously described. Non-limiting examples of suitable arylalkyl groups include benzyl, 2-phenethyl and naphthylmethyl. The bond to the parent moiety is through the alkyl.

30 "Heteroaryl" means a single ring or benzofused heteroaromatic group of 5 to 10 atoms comprised of 1 to 9 carbon atoms and 1 or more heteroatoms independently selected from the group consisting of N, O and S. N-oxides of the ring nitrogens are also included, as well as compounds wherein a ring nitrogen is substituted by a C₁-C₆ alkyl group to form a quaternary amine. Examples of single-ring heteroaryl groups are pyridyl, oxazolyl, isoxazolyl, oxadiazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl,

tetrazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyrazinyl, pyrimidyl, pyridazinyl and triazolyl. Examples of benzofused heteroaryl groups are indolyl, quinolyl, isoquinolyl, phthalazinyl, benzothienyl (i.e., thionaphthenyl), benzimidazolyl, benzofuranyl, benzoxazolyl and benzofurazanyl. All positional isomers are contemplated, e.g., 2-pyridyl, 3-pyridyl and 4-pyridyl.

"Alkoxy" means an alkyl radical attached by an oxygen, i.e., alkoxy groups having 1 to 9 carbon atoms.

"Alkenyl" means straight or branched chains of carbon atoms having one or more double bonds in the chain, conjugated or unconjugated.

"Oxime" means a $\text{CH}(\text{:NOH})$ radical containing moiety.

"Halogen", "halogenated" or "halo" refers to fluorine, chlorine, bromine or iodine radicals.

The term "prodrug," as used herein, represents compounds which are rapidly transformed in vivo to the parent compound of the above formula, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Prodrugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

Linker groups such as L^1 , L^2 , Y and Z are divalent:

In a preferred group of compounds of formula I,



L^1 is $-\text{SO}_2-$, $-\text{CH}_2-$, $-\text{CHCH}_3-$, $-\text{C}(=\text{O})-$, $-\text{C}=\text{NOR}^2-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{CHOH}-$, $-\text{O}-$, $-\text{S}-$ or $-\text{S}=\text{O}$;

L^2 is $-\text{SO}_2-$, $-\text{C}(=\text{O})-$, $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{C}(\text{CH}_3)=\text{CH}_2-$, $-\text{NH}-$, $-\text{O}-$, $-\text{NHSO}_2-$, $-\text{NHC}(\text{O})-$ or $-\text{C}(\text{CH}_3)(\text{OH})-$;

R^1 is H, $-\text{CH}_3\text{NH}_2$, $-\text{CH}_2\text{CF}_3$, $-\text{NHC}_3\text{H}_7$, $-\text{NHC}_2\text{H}_5$, $-\text{NHC}_4\text{H}_9$, C_1-C_6 alkyl, $-\text{CF}_3$, $-\text{CH}(\text{CH}_3)_2$, thiophenyl, morpholinyl, cyclopropanyl, benzyl, naphthyl, $\text{C}(\text{CH}_3)_3$, NHphenyl, 3,5-difluorophenyl, phenyl, N-cyclopentyl or $\text{N}(\text{CH}_3)_2$;


R^2 is H or CH_3 ;

R^4 is furanyl, pyridyl, pyrimidyl, thiophenyl, quinolyl, t-butoxy, alkoxy, cyclohexyl, phenyl, tolyl, C_3H_7 , dimethylpyrimidyl, trifluoromethoxyphenyl, morpholinylphenyl or CH_3 ; with the proviso that when R^4 is t-butoxy, L^2 must be

5  , $-CH_2-$, $-CHCH_3-$, $-C(CH_3)_2-$ or  , all of the above optionally substituted with one to three substituents, which are the same or different and are independently selected from X;

R^5 and R^6 are independently H or CH_3 ;

X is H, Cl, CF_3 , OCH_3 , OCF_3 , OCF_2H , CH_3 or C_1-C_6 cycloalkyl;

10 Y is $-SO_2-$ or  ;

Z is a covalent bond; or

R^1 , Y, R^2 and Z taken together with the nitrogen atom form a morpholinyl group.

15 In a more preferred embodiment of the invention,

L^1 is $-SO_2-$ or $-CH_2-$;

L^2 is $-SO_2-$;

R^1 is CH_3 or CF_3 ; and

20 R^4 is phenyl, pyrimidyl or pyridyl, said phenyl, pyrimidyl or pyridyl groups optionally substituted with one to three substituents independently selected from the group consisting of C_1-C_6 alkyl, C_1-C_6 alkoxy, OH, CF_3 and halogen.

More preferably, the phenyl is substituted with OCH_3 or halogen selected from fluorine and chlorine.

25

Compounds of the invention may have at least one asymmetrical carbon atom and therefore all isomers, including diastereomers and rotational isomers are contemplated as being part of this invention. The invention includes (+)- and (-)- isomers in both pure form and in admixture, including racemic mixtures. Isomers can be prepared using conventional techniques, either by reacting optionally pure or optically enriched starting materials or by separating isomers of a compound of

30

formula I. Those skilled in the art will appreciate that for some compounds of formula I, one isomer may show greater pharmacological activity than other isomers.

Compounds of formula I can exist in unsolvated and solvated forms, including hydrated forms. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol and the like, are equivalent to the unsolvated forms for purposes of this invention.

Compounds of the invention with a basic group can form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salt is prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt. The free base form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium bicarbonate. The free base form differs from its respective salt form somewhat in certain physical properties, such as solubility in polar solvents, but the salt is otherwise equivalent to its respective free base forms for purposes of the invention.

Certain compounds of the invention are acidic (e.g., compounds where R^2 is a hydrogen covalently bonded to N). Acidic compounds according to the present invention can form pharmaceutically acceptable salts with inorganic and organic bases. Examples of such salts are the sodium, potassium, calcium, aluminum, magnesium, zinc, lithium, gold and silver salts. Also included are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine, piperazines and other amines.

Compounds of the present invention are generally prepared by processes known in the art, for example by the processes described below.


The following abbreviations are used in the procedures and schemes: aqueous (aq), anhydrous (anhyd), n-butyllithium (n-BuLi), dibromodimethylhydantoin (DBDMH), diisopropylethylamine (DIPEA), diethyl ether (Et_2O), dimethylacetamide (DMA), dimethyl sulfoxide (DMSO), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), ethanol (EtOH), ethyl acetate (EtOAc), 2-propanol (IPA), leaving group (LG), lithium hexamethyldisilazide (LHMDS), meta-chloroperoxybenzoic acid (MCPBA), methanesulfonic acid (MsOH), methanesulfonyl chloride (MsCl),

N-iodosuccinamide (NIS), preparative thin layer chromatography on Merck- silica plates (PTLC), phenyl (Ph), pyridinium chlorochromate (PCC), pyridine (Py), trifluoroacetic anhydride (TFAA), triflic anhydride (Trf_2O), tetrahydrofuran (THF), silica gel chromatography (sgc), thin layer chromatography (TLC), room temperature (rt), 5 hours (h), minutes (min), molar (M), pounds per square inch (psi), and saturated aqueous sodium chloride solution (brine).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209

Description of Reactions-General Scheme I

In step 1, trifluoroacetic anhydride is dissolved in a suitable inert solvent such as methylene chloride and reacted with a benzyl amine at room temperature for 1-5 hr. MsOH (2 eq) is added followed by DBDMH and the reaction mixture is stirred



overnight at room temperature and subjected to aqueous work up. The crude product is recrystallized from a mixture of Et₂O and hexanes or purified via chromatography.

10 In step 2, the product of step 1 is dissolved in THF, cooled in a dry ice/IPA bath and treated with methyllithium then n-BuLi. The resulting dianion may be trapped with
5 a sulfonyl fluoride or a disulfide. If a disulfide is the trapping agent, the resulting product is oxidized with MCPBA in CH₂Cl₂ at room temperature for 1-6 h. The product may be purified via chromatography or crystallization.

In step 3, the product of step 2 is dissolved in THF and treated with n-BuLi at -78 °C to form a dianion that is trapped with a suitable electrophile.


15 Alternatively, in step 3 the product of step 2 is dissolved in THF treated with n-BuLi at -78 °C to form a dianion which is trapped with iodine to provide the iodo substituted product. The product may be purified via sgc or crystallization. The iodo product can be converted to a similar product by nucleophilic aromatic substitution with a variety of nucleophiles, including amines, alcohols, and thiols.

20 In step 4, the product of step 3 is dissolved in a suitable solvent such as dioxane, ethanol, methanol or THF and an alkali metal hydroxide or carbonate such as lithium hydroxide or potassium carbonate is added either as an aqueous solution or as a solid. The reaction mixture is stirred at room temperature for 0.5-24 h. The product may be purified via sgc or crystallization.

25 In step 5, a combination of the product of step 4 and a tertiary amine base is dissolved in a suitable solvent such as CH₂Cl₂ or dioxane, at room temperature, cooled, and a suitable electrophile is added. The reaction mixture is stirred between -78 °C and 100 °C for 0.5 to 48 h. The product may be purified via sgc or crystallization.

30 In step 6, the product of step 5 is dissolved in a suitable inert solvent such as THF or CH₂Cl₂ and treated with a suitable base such as NaH or triethylamine. An electrophile is added and the reaction mixture is stirred between 0 °C and 100°C for 0.5 to 48 h. The product may be purified via sgc or crystallization.

General Scheme II Preparation of Methylene Linked Compounds



Description of reactions-General Scheme II

- In step 1, trifluoroacetic anhydride is dissolved in a suitable inert solvent such as methylene chloride and treated with a benzyl amine at ambient temperature, then stirred for 1-5 h. Methanesulfonic acid (2 eq) is added followed by dibromodimethylhydantoin and the reaction mixture is stirred overnight at rt and subjected to aqueous work up. The product may be purified by chromatography or crystallization.
- In step 2, the product of step 1 is dissolved in THF, cooled in a dry ice/acetone bath (-78°C) and treated with methylolithium, then n-BuLi. The dianion is then treated with a THF solution containing the aldehyde (i). The resulting mixture is warmed to rt and stirred for 10 h. The product is purified by chromatography.
- In step 3, the alcohol product from step 2 is dissolved in methylene chloride and treated with ten fold excess of triethylsilane followed by a slight excess of boron trifluoride etherate. The resulting mixture is stirred at room temperature for 4h, and purified by chromatography.
- In step 4, the product of step 3 is dissolved in a suitable solvent such as dioxane, ethanol, or THF and an alkali metal hydroxide such as lithium hydroxide is


added either as an aqueous solution or as a solid. The reaction mixture is stirred at rt for 0.5-24 h.

In step 5, the product of step 4 is dissolved in a mixture of a suitable inert solvent such as CH_2Cl_2 or dioxane and a tertiary amine base, and a suitable electrophile is added. The reaction mixture is stirred between -78°C and 100°C for 0.5 to 48 h.

In step 6, the product of step 5 is dissolved in a suitable inert solvent such as THF or CH_2Cl_2 and treated with a suitable base such as NaH or triethylamine. An electrophile is added and the reaction mixture is stirred between 0°C and 100°C for 0.5 to 48 h.

The aldehyde (i) used in step 2 was prepared by one of the following two procedures; 1) Regioselective ortho lithiation of a 4-substituted benzaldehyde, and quenching with a substituted phenyl disulfide followed by oxidation with metachloroperoxybenzoic acid to the sulfone. 2) Base promoted displacement of fluoride from an ortho-fluorobenzaldehyde by a thiophenol, phenol, or aniline.

General Scheme III Preparation of Ketone and Olefin Linked Compounds




Description of reactions-General Scheme III

In step 1 the secondary alcohol, the product of Step 2 in Scheme II is oxidized with PCC, in a suitable inert solvent such as CH_2Cl_2 , to the carbonyl by stirring at rt for 18 h. In step 2, the ketone is treated with the ylide obtained by base treatment of dried methyltriphenylphosphonium bromide, providing the exo methylene product. In

step 3 the trifluoroacetamide group can be hydrolyzed with base and reacted with a variety of acylating, sulfonylating, alkylating and other electrophilic reagents.

The ketone product can be treated with hydroxylamine hydrochloride in pyridine and heated at 80°C for 24 h. The mixture was cooled to room temperature and the solvent removed under reduced pressure. Upon workup and purification, the oxime is obtained.

General Scheme IV Preparation of Oxygen Linked Compounds



Description of reactions-General Scheme IV

- 5 In step 1, 2-bromo-4-chlorophenol and a 4-fluorobenzonitrile are dissolved in a polar aprotic solvent such as DMA in the presence of a suitable base such as potassium hydroxide. The reaction mixture is heated for 0.5-7 days. Preferred temperatures are greater than 60 °C. The reaction mixture is diluted with a suitable extraction solvent such as diethyl ether and washed with water. The solvents are
- 10 removed and the product is purified via sgc.

In step 2, the product of step 1 is dissolved in a solution of diborane in THF. The reaction is stirred at reflux for 1-24 h then quenched with water and partitioned between EtOAc and aq NaOH. The solvents are evaporated and the product is purified by formation of the HCl salt in diethyl ether.

- 5 In step 3, the product of step 2 is suspended in CH_2Cl_2 and a suitable base such as triethylamine is added. The reaction mixture is cooled, and TFAA is added. The reaction mixture is stirred from 0.5 to 8 h, then subjected to aqueous workup. The crude product is purified by sgc.


- 10 In step 4, the product of step 3 is dissolved in THF and treated with methyl lithium, then n-BuLi at -78°C to form a dianion that is trapped with a suitable electrophile. The reaction mixture is quenched with a suitable proton source such as aq NH_4Cl or phosphate buffer then extracted with EtOAc. The product may be purified via sgc or crystallization.

- 15 In step 5, the product of step 4 is dissolved in a suitable solvent such as dioxane, ethanol, or THF and an alkali metal hydroxide such as lithium hydroxide is added either as an aqueous solution or as a solid. The reaction mixture is stirred at rt for 0.5-24 h.

- 20 In step 6, the product of step 5 is dissolved in a mixture of a suitable inert solvent such as CH_2Cl_2 or dioxane and a tertiary amine base, and a suitable electrophile is added. The reaction mixture is stirred between -78°C and 100°C for 0.5 to 48 h.

In step 7, the product of step 6 is dissolved in a suitable inert solvent such as THF or CH_2Cl_2 and treated with a suitable base such as NaH or triethylamine. An electrophile is added and the reaction mixture is stirred between 0°C and 100°C for 0.5 to 48 h.

General Scheme V Preparation of Amide Compounds



Description of Reactions-General Scheme V

In step 1, 1-chloro-4-fluorobenzene is dissolved in anhyd THF and treated with
5 n-BuLi at -78°C to form an anion that is trapped with a suitable electrophile. The product may be purified via sgc or crystallization.


In step 2, the product of step 1 is dissolved in a suitable polar solvent such as acetonitrile or DMA. A benzoic acid containing a nucleophilic moiety such as an OH,
10 NHR, or SH is added, and two or more equivalents of a suitable base such as potassium hydroxide or sodium hydride is added. The reaction mixture may be stirred for 1-24 h at temperatures ranging between 0°C and 150°C . The reaction mixture is partitioned between water and a suitable solvent such as EtOAc. The product may be purified via sgc or crystallization.

In step 3, the product of step 2 is dissolved in CH_2Cl_2 . Pentafluorophenol and
15 EDCI are added. The reaction mixture is stirred at rt for 0.5-24 h then partitioned between water and CH_2Cl_2 . The solvents are evaporated. The product may be purified via sgc or crystallization.

In step 4, the product of step 3 is dissolved in a suitable solvent such as CH_2Cl_2 . An amine base such as DIPEA or triethylamine is added, followed by a
20 primary or secondary amine. The reaction mixture may be stirred for 1-24 h at rt. The reaction mixture is then subjected to aqueous workup and isolation and the product is purified via sgc.

In step 5, if the nucleophilic moiety in step 2 contains oxidizable functionality, the product of step 4 is dissolved in a suitable solvent such as CH_2Cl_2 and MCPBA is added. The reaction mixture may be stirred for 0.5-48 h then partitioned between a suitable solvent such as CH_2Cl_2 or EtOAc and an aqueous base such as Na_2CO_3 . The solvent is evaporated and the product is purified via sgc.

General Scheme VI C-Ring Addition Elimination Chemistry



Description of Reactions-General Scheme VI

In step 1, trifluoroacetic anhydride is dissolved in a suitable inert solvent such as methylene chloride and reacted with a benzyl amine at rt for 1-5 h. Methanesulfonic acid (2 eq) is added followed by N-iodosuccinamide. The reaction mixture is stirred overnight at rt, then subjected to aqueous work up. The crude product is recrystallized from isopropanol and water.

In step 2, CuCl is dissolved in glacial acetic acid. The flask is cooled to 0°C and SO_2 gas is bubbled in with stirring for 40 min. In a separate flask 2-fluoro-4-chloroaniline is dissolved in glacial acetic acid and concentrated HCl. The resulting

solution is cooled to 0 °C and treated with an aqueous solution of NaNO₂. The reaction mixture is stirred for 30 min at 0 °C and the contents are added to the flask containing the SO₂ solution causing vigorous gas evolution. The reaction is then allowed to warm to rt. The product is isolated by pouring the reaction mixture onto chipped ice, then filtering the resulting solid.

In step 3, the product of step 2 is dissolved in acetone. An aqueous solution of KF (2 eq) is added and the reaction mixture is stirred for 12-24 h at rt. The reaction mixture is extracted with a suitable solvent such as CH₂Cl₂ or Et₂O and the solvent is evaporated to afford the product.

In step 4, the product of step 1 is dissolved in THF and TMEDA is added. The flask is placed under N₂ blanket, and cooled to 0 °C. A solution of isopropyl magnesium chloride in THF is added and the reaction mixture is stirred for 1-4 h. The resulting solution is added to a flask containing the product of step 3 that was cooled with an ice-water bath. The reaction mixture is stirred for 1-3 h. The reaction is quenched with aqueous NH₄Cl and extracted with EtOAc. After evaporation of the solvent, the crude product is purified via sgc.

In step 5, the product of step 4 is dissolved in a suitable solvent such as dioxane, ethanol, or THF and an alkali metal hydroxide such as lithium hydroxide is added either as an aqueous solution or as a solid. The reaction mixture is stirred at rt for 0.5-24 h. The product may be purified via sgc or crystallization.

In step 6, the product of step 5 is dissolved in a suitable inert solvent such as CH₂Cl₂ or acetonitrile and a tertiary amine base, and a triflic anhydride is added. The reaction mixture is stirred between -78°C and rt for 0.5 to 48 h. The product may be purified via sgc or crystallization.









In step 7, the product of step 6 is dissolved in a suitable inert solvent such as dioxane and a thiol is added. A base such as sodium hydride, sodium hydroxide, or NaHMDS is added and the reaction mixture is stirred at a suitable temperature between 50 °C and 100 °C for 4-24 h. The reaction mixture is quenched with water and extracted with a suitable solvent. The solvents are evaporated and the crude product is purified via sgc.













In step 8, the product of step 7 is dissolved in a suitable inert solvent such as CH₂Cl₂. Na₂HPO₄ and urea hydrogen peroxide complex is added, followed by TFAA.















The reaction mixture is refluxed for 4-16 h, then partitioned between water and CH_2Cl_2 . The solvents are evaporated and the crude product is purified via sgc.

- Those skilled in the art will appreciate that similar reactions to those described in the above schemes may be carried out on other compounds of formula I as long as substituents present would not be susceptible to the reaction conditions described. Starting materials for the above processes are either commercially available, known in the art, or prepared by procedures well known in the art. Exemplary compounds of formula 1 are set forth below in Table I. CB means covalent bond.
















TABLE I

















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
A	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
B	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
C	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₂ H	SO ₂	CB
D	CH ₃	H	H	t-butoxy	H	CH ₃	SO ₂	CO	OCH ₃	SO ₂	CB
E	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
F	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
G	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CH ₃	SO ₂	CB
H	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
I	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB



















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
J	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₃	SO ₂	CB
K	CH ₃	H	H	t-butoxy	H	CH ₃	SO ₂	C≡O	OCF ₂ H	SO ₂	CB
L	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
M	CH ₃	H	H		H	H	SO ₂	SO ₂	OCH ₃	SO ₂	CB
N	CH ₃	H	H		H	H	CH ₂	SO ₂	OCH ₃	SO ₂	CB
O	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
P	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
Q	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CH ₃	SO ₂	CB
R	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
S	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
T	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
U	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
V	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
W	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
X	CF ₃	H	H		CH ₃	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
Y	C ₆ H ₉	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
Z	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
AA	CH ₃	H	H	C ₃ H ₇	H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
AB	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AC	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AE	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AF	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AG	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AH	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AI	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
AK	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AM	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AO	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
AQ	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AR	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AS	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AT	N(CH ₃) ₂	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AU	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AV	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AW	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AX	CH ₃	H	H	C ₃ H ₇	H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
AY	CF ₃	H	H		CH ₃	CH ₃	SO ₂	SO ₂	Cl	O C	CB
AZ	CF ₃	H	H		CH ₃	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
BA	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₃	SO ₂	CB
BB	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₃	SO ₂	CB
BC	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₃	SO ₂	CB
BD	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₃	SO ₂	CB
















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
BG	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BH	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BJ	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BN	CF ₃	H	H	C ₃ H ₇	H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BO	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BP	NHC ₃ H ₇	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BR	CF ₃	H	H		CH ₃	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BS	CH ₃	H	H		CH ₃	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BT	CH ₃	CH ₃	H		CH ₃	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BU	CH ₃	H	H		H	H	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BV	CH ₃	CH ₃	H		H	H	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BW	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
BX	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BY	CH ₃	CH ₃	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
BZ	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CH ₃	SO ₂	CB
CA	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CH ₃	SO ₂	CB
























	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
CB	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	CH ₃	SO ₂	CB
CC	CF ₃	H	H	C ₃ H ₇	H	CH ₃	SO ₂	SO ₂	Cl	C=O	CB
CD	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
CE	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
CF	-CH(CH ₃) ₂	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CG	NH ₂	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CH	C ₆ H ₅	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CI	-CHCF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CJ		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CK		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CL		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CM		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
CN	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
















	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
CO		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CP	C ₃ H ₇	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CQ		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CR		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CS	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	C=O	CB
CT	NH-(CH ₂) ₂ -CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CU		H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB
CV	CF ₃	H	H		H	CH ₃	SO ₂	C=O	OH	C=O	CB
CW	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OH	SO ₂	CB
CX	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OH	SO ₂	CB
CZ	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₂ H	C=O	CB
DA	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCF ₂ H	SO ₂	CB
DC	CF ₃	H	H		H	CH ₃	SO ₂	C=O	OCH ₃	SO ₂	CB
DD	CF ₃	H	H		H	CH ₃	SO ₂	C=O	OCH ₃	C=O	CB






	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
DE		H	H		H	CH ₃	SO ₂	C=O	OCH ₃	C=O	CB
DF	CH ₃	H	H		H	CH ₃	SO ₂	C=O	Cl	SO ₂	CB
DG	CF ₃	H	H		H	CH ₃	SO ₂	C=O	Cl	SO ₂	CB
DH	CH ₃	H	H		H	CH ₃	SO ₂	CH ₂	Cl	SO ₂	CB
DI	CH ₃	H	H		H	CH ₃	SO ₂	C=O	Cl	SO ₂	CB
DJ	CH ₃	H	H		H	CH ₃	SO ₂	CH ₂	Cl	SO ₂	CB
DK	CH ₃	H	H		H	CH ₃	SO ₂	C=O	Cl	SO ₂	CB
DL	CH ₃	H	H		H	CH ₃	SO ₂	C=CH ₂	Cl	SO ₂	CB
DM	CF ₃	H	H		H	CH ₃	SO ₂		Cl	C=O	CB
DN	CF ₃	H	H		H	CH ₃	SO ₂	C=O	Cl	C=O	CB
DP	CF ₃	CH ₃	H		H	CH ₃	SO ₂	C=CH ₂	Cl	C=O	CB
DQ	CH ₃	H	H		CH ₃	CH ₃	SO ₂	C=O	Cl	SO ₂	CB
DR	CH ₃	H	H		H	CH ₃	SO ₂	NH	Cl	SO ₂	CB
DS	CF ₃	H	H		H	CH ₃	SO ₂	O	Cl	C=O	CB

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
DU	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCH ₃	SO ₂	CB
DV	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	Cl	SO ₂	CB
DW	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
DX	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
DY	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
DZ	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
EA	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
EC	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	CF ₃	SO ₂	CB
ED	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	C=O	CB
EE	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EG	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EH	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EI	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EJ	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
EK	CF ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EL	CH ₃	H	H		H	H	CH ₂	SO ₂	OCH ₃	SO ₂	CB
EN	CF ₃	H	H		CH ₃	CH ₃	CH ₂	SO ₂	OCF ₃	C=O	CB
EP	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
EU	CH ₃	H	H		H	CH ₃	C=O	SO ₂	OCF ₃	SO ₂	CB
EV	CF ₃	H	H		H	CH ₃	C=O	SO ₂	OCH ₃	SO ₂	CB
EW	CF ₃	H	H		H	CH ₃	C=O	O	H	C=O	CB
EX	CF ₃	H	H		H	CH ₃	C=O	O	H	C=O	CB
EY	CF ₃	H	H		H	CH ₃	C=O	O	Cl	C=O	CB
EZ	CH ₃	H	H		H	CH ₃	C=O	SO ₂	OCF ₃	C=O	CB
FA	CF ₃	H	H		H	CH ₃	C=O	NHSO ₂	H	C=O	CB
FB	CF ₃	H	H		H	CH ₃	C=O	NHCO	H	C=O	CB
FC	CF ₃	H	H		H	CH ₃	C=CH ₂	SO ₂	OCF ₃	C=O	CB
FD	CH ₃	H	H		H	CH ₃	C=CH ₂	SO ₂	OCF ₃	SO ₂	CB
FE	CF ₃	H	H		H	CH ₃	C=O	SO ₂	OCF ₃	SO ₂	CB

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
FF	CH ₃	H	H		H	CH ₃	C=NO H	SO ₂	OCF ₃	SO ₂	CB
FG	CH ₃	H	H		H	CH ₃	C(CH ₃) ₂	SO ₂	Cl	SO ₂	CB
FH	CF ₃	H	H		H	H	C=O	SO ₂	OCH ₃	C=O	CB
FI	CH ₃	H	H		H	H	O	SO ₂	Cl	SO ₂	CB
FJ	*R ¹ , Y, Z and R ² combine to form morpholinyl	*	H		-		S	SO ₂	Cl	*	*
FK	H	CH ₃	H		-		S=O	SO ₂	Cl	CB	CB
FL	H	CH ₃	H		-		SO ₂	SO ₂	Cl	CB	CB
FM	*R ¹ , Y, Z and R ² combine to form morpholinyl	*	H		-		SO ₂	SO ₂	Cl	*	*
FN	CH ₃	CH ₃	H		-		S	SO ₂	Cl	CB	CB
FO	H	H	H		-		S	SO ₂	Cl	CB	CB
FP	CH ₃	CH ₃	H		-		SO ₂	SO ₂	Cl	CB	CB
FQ	H	H	H		-		SO ₂	SO ₂	Cl	CB	CB
FR	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
FS	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
FT	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB


	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
FU	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	C=O	CB
FV	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
FW	CH ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
FX	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	C=O	CB
FY	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
FZ	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	H	SO ₂	CB
GA	CH ₃	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	C=O	CB
GD		H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
GF	NHC ₂ H ₅	H	H		H	CH ₃	CH ₂	SO ₂	OCF ₃	SO ₂	CB
GG	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
GH	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	CF ₃	SO ₂	CB
GI	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	Cl	SO ₂	CB
GJ	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	SO ₂	CB
GK	CF ₃	H	H		H	CH ₃	SO ₂	C=O	OCH ₃	SO ₂	CB

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	L ¹	L ²	X	Y	Z
GL	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OH	SO ₂	CB
GM	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH(CH ₃) ₂	SO ₂	CB
GN	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂		SO ₂	CB
GO	CF ₃	H	H		H	CH ₃	SO ₂	SO ₂	OCH ₃	C=O	CB



CB is a covalent bond
- means that the substituent is not present













5


















In a preferred embodiment, there are disclosed compounds of the formula














- 10 a prodrug thereof, or a pharmaceutically acceptable salt of the compound or of said prodrug; wherein X, R¹ and R⁴ are as shown in the table below:


Example	X	R ¹	R ⁴
A	OCH ₃	CH ₃	
C	OCF ₂ H	CH ₃	

Example	X	R ¹	R ⁴
G	CH ₃	CH ₃	
L	Cl	CH ₃	
R	CF ₃	CF ₃	
S	Cl	CF ₃	
AB	CF ₃	CH ₃	
AT	Cl	N(CH ₃) ₂	
BA	OCF ₃	CH ₃	
BD	OCF ₃	CF ₃	
BZ	CH ₃	CF ₃	
CD	Cl	CH ₃	
FS	H	CH ₃	
FY	H	CF ₃	

Example	X	R ¹	R ⁴
GG	Cl	CF ₃	
GH	CF ₃	CF ₃	
XXIX		CF ₃	
XXX		CF ₃	
XXXI		CF ₃	
XXXII	CN	CF ₃	
XXXIII	NH ₂	CF ₃	
XXXIV		CF ₃	
XXXVI	Cl	CF ₃	
XXXVII		CF ₃	
XXXVIII	CN	CF ₃	
XXXIX	-CONH ₂	CF ₃	



















Example	X	R ¹	R ⁴
XXXX	-OCH ₃	CF ₃	
XXXXI	-OH	CF ₃	
XXXXII		CF ₃	
XXXXIII		CF ₃	
XXXXIV	H ₃ C-CH ₂ -O-	CF ₃	
XXXXV	H ₃ C-O-CH ₂ -CH ₂ -O-	CF ₃	
XXXXXV	OCH ₃	CF ₃	
XXXXXVI		CH ₃	

In another preferred embodiment, there are disclosed compounds of the formula



5

a prodrug thereof, or a pharmaceutically acceptable salt of the compound or of said prodrug; wherein X, Y-R¹ and R⁴ are as shown in the table below:

Example	X	Y-R ¹	R ⁴
XXXXVI			
XXXXVII			
XXXXVIII			
XXXXXIX			
XXXXXXI	-OCH ₃		
XXXXXII	-OCH ₃		
XXXXXIII	-OCH ₃		

Compound A : ¹H NMR (300 MHz, CDCl₃) 1.54 (d, J = 6.9Hz 3H), 2.67 (s, 3H), 4.72 (q, J = 5Hz 1H), 4.86 (br. d, J = 5Hz, 1H, NH), 7.08-8.42 (m, 11H).

- 5 Compound C : ¹H NMR (400 MHz, CDCl₃) 1.51 (d, J = 7.2Hz 3H), 2.67 (s, 3H), 4.702 (q, J = 6.8Hz 1H), 5.05 (br. d, J = 6.4Hz, 1H, NH), 6.71 (t, J = 71.6 Hz, CF₂H) 7.07-8.47 (m, 11H).

- 10 Compound G : ¹H NMR (300 MHz, CDCl₃) 8.43-8.41 (m, 1H), 8.36 (d, 8Hz, 1H), 8.28-8.22 (m, 1H), 7.96-7.92 (m, 2H), 7.69-7.60 (m, 2H), 7.52-7.47 (m, 2H), 7.43-7.37 (m, 1H), 7.13-7.06 (m, 1H), 4.76-4.70 (m, 2H), 2.68 (s, 3H), 2.59 (s, 3H), 1.41 (d, 7 Hz, 3H).

- 15 Compound L : ¹H NMR (300 MHz, CDCl₃) 8.61-5.97 (m, 2H), 8.40 (d, 8 Hz, 1H), 8.24-8.21 (m, 1H), 7.96 (d, 8 Hz, 2H), 7.86-7.83 (m, 1H), 7.70-7.63 (m, 1H), 7.52 (d, 8

Hz, 2H), 7.46-7.40 (m, 1H), 7.18-7.12 (m, 1H), 4.80-4.70 (m, 1H), 2.71 (s, 3H), 1.56 (d, 7Hz, 3H).

5 Compound R : ^1H NMR (300 MHz, CDCl_3) 8.89-8.87 (m, 1H), 8.58 (d, 8Hz, 1H), 8.32-8.25 (m, 1H), 8.15-8.11 (m, 1H), 8.03-7.98 (m, 2H), 7.71-7.63 (m, 1H), 7.52-7.48 (m, 2H), 7.47-7.41 (m, 1H), 7.16-7.09 (m, 1H), 5.62 (d, 8 Hz, 1H), 4.90-4.80 (m, 1H), 1.63 (d, 7 Hz, 3H).

10 Compound S : ^1H NMR (300 MHz, CDCl_3) 8.61-8.59 (m, 1H), 8.39 (d, 8 Hz, 1H), 8.29-8.24 (m, 1H), 7.99 (d, 8 Hz, 2H), 7.86-7.82 (m, 1H), 7.67-7.62 (m, 1H), 7.49 (d, 8Hz, 1H), 7.46-7.40 (m, 1H), 7.16-7.10 (m, 1H), 4.89-4.84 (m, 1H), 1.65 (d, 6 Hz, 1H).

15 Compound AB : ^1H NMR (300 MHz, CDCl_3) 8.88-8.86 (m, 1H), 8.62-8.59 (m, 1H), 8.30-8.29 (m, 1H), 8.15-8.11 (m, 1H), 8.00-7.96 (m, 2H), 7.71-7.63 (m, 1H), 7.56-7.52 (m, 2H), 7.47-7.41 (m, 1H), 7.16-7.09 (m, 1H), 4.99-4.84 (m, 1H), 4.80-4.70 (m, 1H), 2.71 (s, 3H), 1.54 (d, 7Hz, 3H).

20 Compound AT : ^1H NMR (300 MHz, CDCl_3) 8.51 (br s 1H), 8.39 (d, 8 Hz, 2H), 7.99 (d, 8 Hz, 2H), 7.86-7.83 (m, 1H), 7.61-7.50 (m, 1H), 7.49 (d, 8 Hz), 7.05-6.99 (m, 1H), 4.70-4.50 (m, 2H), 2.83 (s, 3H), 2.57 (s, 3H), 1.50 (d, 7 Hz, 3H).

Compound BA : ^1H NMR (300 MHz, CDCl_3) 1.54 (d, J = 6.9 Hz 3H), 2.7 (s, 3H), 4.72 (q, J = 5.5Hz 1H), 5.05 (br. d, J = 5Hz, 1H, NH), 7.1 -8.55 (m, 11H).

25 Compound BD : ^1H NMR (300 MHz, CDCl_3) 8.51 (d, 9 Hz, 1H), 8.47-8.45 (m, 1H), 8.01-7.97 (m, 2H), 7.71-7.63 (m, 2H), 7.52-7.41 (m, 3H), 7.17-7.10 (m, 1H), 5.51 (d, 8 Hz, 1H), 4.90-4.80 (m, 1H), 1.64 (d, 7 Hz, 3H).

30 Compound BZ : ^1H NMR (300 MHz, CDCl_3) 8.43 (br s, 1H), 8.32 (d, 8 Hz, 1H), 8.28-8.22 (m, 1H), 7.94 (d, 8 Hz, 2H), 7.68-7.58 (m, 2H), 7.47-7.37 (m, 3H), 7.12-7.06 (m, 1H), 5.72 (d, 8 Hz, 1H), 4.86-4.70 (m, 1H), 2.59 (s, 3H), 1.60 (d, 7 Hz, 3H).

35 Compound CD : ^1H NMR (300 MHz, CDCl_3): 8.82-8.78 (m, 1H), 8.23 (d, 7 Hz, 2H), 8.21-8.07 (m, 1H), 7.81-7.77 (m, 2H), 7.63-7.57 (m, 1H), 7.55 (d, 7 Hz, 2H), 7.40-7.32 (m, 1H), 7.20-7.16 (m, 1H), 4.8-4.7 (m, 2H), 2.67 (s, 3H), 1.55 (d, 7 Hz, 2H).

40 Compound FS : ^1H NMR (300 MHz, CDCl_3) 8.66-8.62 (m, 1H), 8.51-8.47 (m, 1H), 8.29-8.24 (m, 1H), 7.99-7.95 (m, 2H), 7.93-7.89 (m, 2H), 7.67-7.53 (m, 1H), 7.50-7.44 (m, 2H), 7.42-7.39 (m, 1H), 7.13-7.07 (m, 1H), 4.78-4.73 (m, 1H), 4.61-4.59 (m, 1H), 2.70 (s, 3H), 1.56 (d, 7 Hz, 3H).

45 Compound FY : ^1H NMR (300 MHz, CDCl_3) 8.66-8.63 (m, 1H), 8.49-8.46 (m, 1H), 8.28-8.25 (m, 1H), 8.01 (d, 8 Hz, 2H), 7.93-7.89 (m, 2H), 7.65-7.58 (m, 1H), 7.56 (d, 8 Hz, 2H), 7.47-7.41 (m, 1H), 7.13-7.07 (m, 1H), 5.18 (d, 6 Hz, 1H), 4.90-4.80 (m, 1H), 1.66 (d, 7 Hz, 3H).

Compound GG : ^1H NMR (300 MHz, CDCl_3): 8.88 (d, 1.2 Hz, 1H), 8.51-8.56 (m, 2H), 8.31 (dd, 8 Hz, 1Hz, 1H), 8.18 (dd, 8 Hz, 1 Hz, 1H), 8.08-7.96 (m, 3H), 7.62-7.48 (m, 3H), 5.51 (d, 9 Hz, 1H), 4.90-4.70 (m, 1H), 1.62 (d, 7 Hz, 3H).

5 Compound GH : ^1H NMR (300 MHz, CDCl_3): 8.63 (d, 2 Hz), 8.58-8.55 (m, 1H), 8.34-8.28 (m, 2H), 8.07-7.98 (m, 3H), 8.35 (dd, 8 Hz, 2 Hz, 1H), 7.55-7.46 (m, 3H), 5.34 (d, 8 Hz, 1H), 4.9-4.8 (m, 1H), 1.64 (d, 6 Hz, 3H).

10 Compound GQ/XXIX, : ^1H NMR (300 MHz, CDCl_3): δ 8.56-8.52 (m, 1H), 8.32-8.21 (m, 3H), 8.02-7.92 (m, 4H), 5.42 (d, 9 Hz, 1H), 8.02-7.92 (m, 4H), 5.42 (d, 1H, 9 Hz), 4.84-4.78 (m, 1H), 2.16-2.06 (m, 1H), 1.60 (d, 7Hz, 3H), 1.20-1.17 (m, 2H), 0.97-0.89 (m, 1H).

15 Compound GR/XXX, : ^1H NMR (300 MHz, CDCl_3): δ 8.33-8.22 (m, 3H), 8.00-7.94 (m, 2H), 7.66-7.58 (m, 1H), 7.53-7.37 (m, 4H), 7.16-7.05 (m, 1H), 5.160 (d, 9 Hz, 1H), 4.88-4.83 (m, 1H), 2.17-2.06 (m, 1H), 1.65 (d, 7 Hz, 3H), 1.28-1.20 (m, 2H), 0.97-0.90 (m, 2H).

20 Compound GS/XXXI, : ^1H NMR (300 MHz, CDCl_3): δ 8.38-8.29 (m, 2H), 8.17 (d, 8 Hz, 1H), 8.07-8.02 (m, 1H), 7.91-7.85 (m, 2H), 7.56-7.36 (m, 5H), 6.11 (d, 8 Hz, 1H), 4.84-4.78 (m, 1H), 2.12-2.01 (m, 1H), 1.57 (d, 7Hz, 3H), 1.21-1.12 (m, 2H), 0.92-0.86 (m, 2H).

25 Compound GW/XXXVI, : ^1H NMR (300 MHz, CDCl_3): δ 10.19 (d, 7.8 Hz, 1H), 8.27-8.42 (m, 4H), 8.13 (dd, 7.8 Hz, 2.1 Hz, 1H), 7.93 (d, 8.4 Hz, 2H), 7.78-7.63 (m, 2H), 7.59 (d, 8.4 Hz, 2H), 4.80 (m, 1H), 1.44 (d, 6.9 Hz, 3H).

30 Compound HO/XXXXXV, : ^1H NMR (300 MHz, CDCl_3): δ 8.56 (d, 3.9 Hz, 1H), 8.31-8.22 (m, 2H), 8.124 (d, 2.7 Hz, 1H), 8.05-7.95 (m, 1H), 7.92 (d, 8.4 Hz, 2H), 7.50-7.45 (m, 1H), 7.92 (d, 8.4 Hz, 2H), 7.27-7.23 (m, 2H), 5.8 (d, NH, 1H), 4.85-4.75 (m, 1H), 3.99 (s, 3H), 1.58 (d, 7.2 Hz, 3H).

35 Compound HP/XXXXXVI, : ^1H NMR (300 MHz, CDCl_3): δ 8.56-8.52 (m, 1H), 8.31-8.23 (m, 3H), 8.02-7.90 (m, 4H), 4.87-4.78 (d, 7 Hz, 1H), 4.69 (m, 1 H), 2.66 (s, 3H), 2.16-2.06 (m, 1H), 1.51 (d, 7 Hz, 3H), 1.27-1.17 (m, 2H), 0.96-0.90 (m, 2H).

The compounds of the present invention exhibit anti-inflammatory and/or immunomodulatory activity and are useful in the treatment of various medical

40 conditions including, e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, osteoporosis, renal ischemia, cerebral stroke, cerebral ischemia, nephritis, psoriasis, allergy, inflammatory disorders of the lungs and gastrointestinal tract such as Crohn's disease, and respiratory tract disorders such as reversible airway obstruction, asthma, chronic obstructive pulmonary disease (COPD)

and bronchitis. This utility is manifested as demonstrated by activity in the following assay.

Potential cannabinoid receptor ligands were screened for the ability to compete with [3 H] CP-55,940 for binding to recombinant cannabinoid receptors. Test

- 5 compounds were serially diluted in Diluent Buffer (50 mM Tris pH 7.1, 1 mM EDTA, 3 mM $MgCl_2$, 0.1% BSA, 10% DMSO, 0.36% methyl cellulose (Sigma M-6385)) from stocks prepared in 100% DMSO. Aliquots (10 μ l) were transferred into 96-well microtiter plates. Membrane preparations of recombinant human cannabinoid CB2 receptor (Receptor Biology #RB-HCB2) or recombinant human cannabinoid CB1
- 10 receptor (Receptor Biology #RB-HCB1) were diluted to 0.3 mg/ml in Binding Buffer (50 mM Tris pH 7.2, 1 mM EDTA, 3 mM $MgCl_2$, 0.1% BSA). Aliquots (50 μ l) were added to each well of the microtiter plate. The binding reactions were initiated by addition of [3 H] CP-55,940 (New England Nuclear # NET 1051; specific activity =180 Ci/mmol) to each well of the microtiter plate. Each 100 μ l reaction mixture contained
- 15 0.48 nM [3 H] CP-55,940, 15 μ g membrane protein in binding buffer containing 1% DMSO and 0.036 % methyl cellulose. Following incubation for 2 hours at room temperature, the reactions were filtered through 0.5% polyethylenimine-coated GF/C filter plates (UniFilter-96, Packard) with a TomTec Mark 3U Harvester (Hamden, CT). The filter plate was washed 5 times with binding buffer, rotated 180°, then re-washed
- 20 5 times with binding buffer. Bound radioactivity was quantitated following addition of 30 μ l of Packard Microscint 20 scintillant in a Packard TopCount NXT microplate scintillation counter. Non-linear regression analysis of the resulting data was performed using Prism 2.0b (GraphPad, San Diego, CA).

- Cannabinoid receptor ligands according to the present invention have anti-
- 25 inflammatory activity and/or immunomodulatory activity and are useful in the treatment of various medical conditions including, e.g., cutaneous T cell lymphoma, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, glaucoma, diabetes, osteoporosis, renal ischemia, myocardial infarction, cerebral stroke, cerebral ischemia, nephritis, hepatitis, glomerulonephritis, cryptogenic fibrosing aveolitis,
- 30 psoriasis, atopic dermatitis, vasculitis, allergy, seasonal allergic rhinitis, Crohn's disease, inflammatory bowel disease, reversible airway obstruction, adult respiratory distress syndrome, asthma, chronic obstructive pulmonary disease (COPD) or

bronchitis. It is contemplated that a compound of this invention may be useful in treating more than one of the diseases listed.

Additionally, a compound of the present invention may be co-administered or used in combination with disease-modifying antirheumatic drugs (DMARDs) such as methotrexate, azathioprine, leflunomide, pencillinamine, gold salts, mycophenolate mofetil, cyclophosphamide and other similar drugs. They may also be co-administered with or used in combination with NSAIDs such as piroxicam, naproxen, indomethacin, ibuprofen and the like; COX-2 selective inhibitors such as Vioxx® and Celebrex®; COX-1 inhibitors such as Feldene; immunosuppressives such as steroids, cyclosporine, Tacrolimus, rapamycin and the like; biological response modifiers (BRMs) such as Enbrel, Remicade, IL-1 antagonists, anti-CD40, anti-CD28, IL-10, anti-adhesion molecules and the like; and other anti-inflammatory agents such as p38 kinase inhibitors, PDE4 inhibitors, TACE inhibitors, chemokine receptor antagonists, Thalidomide and other small molecule inhibitors of pro-inflammatory cytokine production. Other drugs that the compounds of the invention may be co-administered or used in combination with include Anaprox, Arava, Arthrotec, Azulfidine, Aspirin, Cataflam, Celestone Soluspan, Clinoril, Cortone Acetate, Cuprimine, Daypro, Decadron, Depen, Depo-Medrol, Disalcid, Dolobid, Naprosyn, Gengraf, Hydrocortone, Imuran, Indocin, Lodine, Motrin, Myochrysine, Nalfon, Naprelan, Neoral, Orudis, Oruvail, Padiapred, Plaquenil, Prelone, Relafen, Solu-Medrol, Tolectin, Trilisate and Volataren. These include any formulation of the abovenamed drugs.

For the treatment of multiple sclerosis, the compounds of the invention may be co-administered or used in combination with Avonex, Betaseron and Copaxone.

For combination treatment with more than one active agent, where the active agents are in separate dosage formulations, the active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of the other agent.

The present invention also relates to a pharmaceutical composition comprising a compound of formula I of this invention and a pharmaceutically acceptable carrier.

The compounds of formula I can be administered in any conventional dosage form known to those skilled in the art. Pharmaceutical compositions containing the compounds of formula I can be prepared using conventional pharmaceutically acceptable excipients and additives and conventional techniques. Such

pharmaceutically acceptable excipients and additives include non-toxic compatible fillers, binders, disintegrants, buffers, preservatives, anti-oxidants, lubricants, flavorings, thickeners, coloring agents, emulsifiers and the like. All routes of administration are contemplated including, but not limited to, parenteral, transdermal, subcutaneous, intramuscular, sublingual, inhalation, rectal and topical.

Thus, appropriate unit forms of administration include oral forms such as tablets, capsules, powders, cachets, granules and solutions or suspensions, sublingual and buccal forms of administration, aerosols, implants, subcutaneous, intramuscular, intravenous, intranasal, intraocular or rectal forms of administration.

When a solid composition is prepared in the form of tablets, e.g., a wetting agent such as sodium lauryl sulfate can be added to micronized or non-micronized compounds of formula I and mixed with a pharmaceutical vehicle such as silica, gelatin starch, lactose, magnesium stearate, talc, gum arabic or the like. The tablets can be coated with sucrose, various polymers, or other appropriate substances.

Tablets can be treated so as to have a prolonged or delayed activity and so as to release a predetermined amount of active principle continuously or at predetermined intervals, e.g., by using ionic resins and the like.

A preparation in the form of gelatin capsules may be obtained, e.g., by mixing the active principle with a diluent, such as a glycol or a glycerol ester, and incorporating the resulting mixture into soft or hard gelatin capsules.

A preparation in the form of a syrup or elixir can contain the active principle together, e.g., with a sweetener, methylparaben and propylparaben as antiseptics, flavoring agents and an appropriate color.

Water-dispersible powders or granules can contain the active principle mixed, e.g., with dispersants, wetting agents or suspending agents, such as polyvinylpyrrolidone, as well as with sweeteners and/or other flavoring agents.

Rectal administration may be provided by using suppositories which may be prepared, e.g., with binders melting at the rectal temperature, for example cocoa butter or polyethylene glycols.

Parenteral, intranasal or intraocular administration may be provided by using, e.g., aqueous suspensions, isotonic saline solutions or sterile and injectable solutions containing pharmacologically compatible dispersants and/or solubilizers, for example, propylene glycol or polyethylene glycol.

Thus, to prepare an aqueous solution for intravenous injection, it is possible to use a co-solvent, e.g., an alcohol such as ethanol or a glycol such as polyethylene glycol or propylene glycol, and a hydrophilic surfactant such as Tween® 80. An oily solution injectable intramuscularly can be prepared, e.g., by solubilizing the active principle with a triglyceride or a glycerol ester.

Topical administration can be provided by using, e.g., creams, ointments or gels.

Transdermal administration can be provided by using patches in the form of a multilaminate, or with a reservoir, containing the active principle and an appropriate solvent.


Administration by inhalation can be provided by using, e.g., an aerosol containing sorbitan trioleate or oleic acid, for example, together with trichlorofluoromethane, dichlorofluoromethane, dichlorotetrafluoroethane or any other biologically compatible propellant gas; it is also possible to use a system containing the active principle, by itself or associated with an excipient, in powder form.

The active principle can also be formulated as microcapsules or microspheres, e.g., liposomes, optionally with one or more carriers or additives.

Implants are among the prolonged release forms which can be used in the case of chronic treatments. They can be prepared in the form of an oily suspension or in the form of a suspension of microspheres in an isotonic medium.


The daily dose of a compound of formula I for treatment of a disease or condition cited above is about 0.001 to about 100 mg/kg of body weight per day, preferably about 0.001 to about 10 mg/kg. For an average body weight of 70 kg, the dosage level is therefore from about 0.1 to about 700 mg of drug per day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

EXAMPLE 1




Compound 1

Compound 1. TFAA (67 mL, 0.474 mol) was dissolved in CH_2Cl_2 (300 mL) and cooled in an ice water bath. A solution of (S)- α -methylbenzylamine (56.4 g, 0.465 mol) dissolved in CH_2Cl_2 (100 mL) was added and the ice bath was removed. The reaction mixture was stirred at rt for 3 h. The reaction mixture was cooled in an ice bath and MsOH (80 mL, 1.23 mol) was added followed by DBDMH (65 g, 0.227 mol). The reaction mixture was left stirring overnight at rt then quenched with 1M aq NaHSO_3 . The organic layer was washed with water and brine, dried with MgSO_4 , and concentrated to give 130 g of white solid. The crude product was recrystallized from Et_2O and hexanes giving 46 g (32%) of intermediate Compound 1 as a solid.



Compound 2

Compound 2. In a flame dried flask under N_2 blanket, Compound 1 (12.35 g, 41.2 mmol) was dissolved in dry THF (165 mL) and cooled to -78°C . Methylolithium (1.4 M in Et_2O , 30 mL, 42 mmol) was added and the reaction mixture was stirred for 5 min. $n\text{-BuLi}$ (1.6 M in hexanes, 26 mL, 42 mmol) was added followed after 10 min by p-methoxybenzenesulfonyl fluoride (8.64 g, 45.4 mmol) which was prepared by standard methods. The cold bath was removed after 10 min and the reaction mixture was allowed to warm to rt over 45 min then quenched with pH 7 sodium phosphate buffer (1 M, 100 mL, 100 mmol). The reaction mixture was extracted with EtOAc and the resulting organic layer was washed with brine and dried with MgSO_4 . After evaporation of the solvent, the crude product was purified by sgc (20%-50% EtOAc /hexanes gradient) to give 10.39 g (65%) of Compound 2 as a solid.




5

Compound 3


Compound 3. In a flame dried flask under N₂ blanket, Compound 2 (11.09 g, 28.6 mmol) was dissolved in anhyd THF (100 mL) and cooled to -78°C. A solution of n-BuLi (2.5 M in hexanes, 24 mL, 60 mmol) was added and the reaction mixture was stirred for 40 min. Bis-4-methoxyphenyl disulfide (8.76 g / 31.5 mmol) was added and the reaction mixture was stirred at -78 °C for 40 min then between -15 °C and -30 °C for 5 h then quenched with pH 7.0 sodium phosphate buffer (1.0 M, 120 mL). The reaction mixture was partitioned between EtOAc and water. The aqueous layer was extracted with additional EtOAc. The combined organic layer was washed with aq Na₂CO₃ and brine, then dried with MgSO₄ and concentrated to dryness. The crude product (13.8 g yellow foam) was dissolved in CH₂Cl₂ (120 mL) and cooled to 0°C. MCPBA (18.5 g, ca 107 mmol) was added, followed by additional CH₂Cl₂ (40 mL). The ice bath was removed and the reaction mixture was stirred at rt for overnight. Aqueous NaHCO₃ (200 mL) and CH₂Cl₂ were added and the layers were separated. The organic layer was washed with aq NaHSO₃, NaHCO₃, H₂O, and brine then dried with MgSO₄. The crude product was purified by sgc (30% to 50% EtOAc/hexanes gradient) to give 7.21 g (45%) of Compound 3.

25




Compound 4

Compound 4. Compound 3 (4.47 g, 8.02 mmol) was dissolved in p-dioxane (16 mL) and cooled to 0 °C. LiOH (1.0 M aq, 10 mL, 10 mmol) was added and the ice bath was removed. The reaction mixture was stirred at rt for 6 h then concentrated. CH₂Cl₂ (100 mL) and NaOH (1.0 M aq, 10 mL) were added and the layers were
 5 separated. The aqueous layer was extracted with additional CH₂Cl₂ and the combined organic layer was dried with MgSO₄ and concentrated. The crude product was purified by sgc (2%-10% MeOH (NH₃)/CH₂Cl₂ gradient mobile phase) to give 3.23 g (87%) of Compound 4.




Compound I

Compound I. Compound 4 (3.08 g, 6.67 mmol) was dissolved in CH₂Cl₂ (33 mL) and triethylamine (1.40 mL, 10.0 mmol) then cooled to 0 °C. MsCl (569 μL, 7.34 mmol) was added and the reaction mixture was stirred at 0 °C for 1 h and 15 min. Citric acid (0.5 M, 40 mL) and additional CH₂Cl₂ were added and the
 15 layers were separated. The organic layer was washed with citric acid, NaHCO₃, and
 20 brine then dried with MgSO₄. The solvent was evaporated and the crude product was purified by sgc (40%-70% EtOAc/hexanes gradient) to give 3.44 g (96%) of Compound I as a solid.



Compound II


Compound II. Compound 4 (27.5 mg, 0.0595 mmol) was dissolved in methylene chloride (226 μ L) and DIPEA (12 μ L). A solution of propionyl chloride dissolved in 1,2-dichloroethane (1 M, 75 μ L, 0.075 mmol) was added and the reaction mixture was shaken at room temperature overnight. Tris(2-aminoethyl)amine
 5 polystyrene (4.1 mmol N/g, ca 60 mg) was added to the reaction mixture. The reaction mixture was shaken for an additional hour at rt. The crude product was concentrated, then dissolved in EtOAc and filtered through a silica-gel SepPak (Waters Corp.). The resulting filtrate was concentrated to give 9 mg (29%) of Compound II.



Compound III

Compound III. Compound 4 (25 mg, 0.054 mmol) was dissolved in CH_2Cl_2 (270 μ L). A solution of phenyl isocyanate dissolved in toluene (1.0 M, 65 mL, 0.065 mmol) was added and the reaction mixture was shaken at rt overnight. Tris(2-aminoethyl) amine polystyrene (4.1 mmol N/g, ca 60 mg) was added to the reaction mixture and the reaction mixture was shaken for an additional 40 min at rt. EtOAc
 20 was added and the reaction mixture was filtered through a silica gel SepPak (Waters Corp.). The resulting filtrate was concentrated to give 18 mg (57%) of Compound III.


EXAMPLE II



Compound 5

Compound 5. In a 3-necked flame-dried flask under N_2 blanket Compound 1 (40.0 g, 134 mmol) was dissolved in anhyd THF (535 mL) and cooled to -75°C


(internal temperature). A solution of methyllithium (1.4 M in diethyl ether, 105 mL, 147 mmol) was added at a rate that kept the internal temperature below -60 °C. The reaction was stirred for 15 min and a solution of n-BuLi (2.5 M in hexanes, 62 mL, 155 mmol) was added at a rate that maintained the internal temperature of the reaction below -65 °C. The reaction mixture was stirred for 40 min. and a solution of bis(4-chlorophenyl) disulfide (42 g, 146 mmol) dissolved in anhyd THF (90 mL) was added via addition funnel over 1 h. The reaction mixture was stirred for 3 h then quenched with HCl (1 M aqueous, 200 mL, 200 mmol). EtOAc (500 mL) was added and the layers were separated. The aqueous layer was extracted with 500 mL EtOAc, and the combined organic layer was washed with 1 M aq KOH, water, and brine. After drying with MgSO₄, the solvent was evaporated to give 54.1 g of a solid. The crude product (52.3 g) was dissolved in CH₂Cl₂ (750 mL) and cooled to 2 °C (internal temp). MCPBA (60%, 184 g) was added in portions over 1 hr and 20 min keeping the internal temperature below 15 °C. The reaction mixture was stirred an additional 2 h. NaOH (1 M aq, 500 mL) and CH₂Cl₂ were added and the layers were separated. The aqueous layer was extracted with an additional 300 mL of CH₂Cl₂. The combined organic layer was washed with 1M aqueous NaOH, water, and brine, then dried with MgSO₄. After evaporation of the solvent, a solid (65 g) was obtained. The crude product was partially purified by trituration from Et₂O/hexanes to give 33.3 g of a solid which was subsequently purified via sgc (20%-25% EtOAc/hexanes) to give 30 g (57%) of Compound 5 as a solid.



Compound 6


Compound 6. In a flame dried 3-necked flask under N₂ blanket Compound 5 (44 g, 112 mmol) was dissolved in anhyd THF (450 mL) and cooled in a dry ice/IPA bath. A solution of n-butyl lithium (2.5 M in hexanes, 92 mL, 230 mmol) was added at a rate that maintained the internal reaction temperature below -60 °C, and the reaction mixture was stirred for 1 h. A solution of 2-fluorobenzenesulfonyl fluoride (22.3 g, 125 mmol) dissolved in anhyd THF (20 mL) was added and the reaction

mixture was left stirring overnight and allowed to warm to rt. The reaction mixture was cooled to 0 °C and quenched with saturated aq ammonium chloride (300 mL). EtOAc (600 mL) and brine (25 mL) were added and the layers were separated. The organic layer was washed with water and brine, then dried with MgSO₄. The solvents were
 5 evaporated giving a foam (62 g). The product was purified by sgc (20%-25% EtOAc/hexanes mobile phase) giving 9.1 g (15%) of Compound 6.



Compound 7

Compound 7. Compound 6 (6.77 g, 12.3 mmol) was dissolved in dioxane (15 mL) and cooled in an ice bath. Aqueous lithium hydroxide (1 M, 15 mL, 15 mmol) was added and the reaction mixture was left stirring overnight. The reaction mixture was concentrated, then partitioned between CH₂Cl₂ and water. The aqueous layer was
 15 extracted with additional CH₂Cl₂ and the combined organic layer was dried with MgSO₄. Evaporation of the solvent afforded 5.66 g of a foam which was purified by sgc (10% MeOH (NH₃)/CH₂Cl₂) to give 4.27 g of Compound 7 (77%).




Compound IV

Compound IV. Compound 7 (2.66 g, 5.86 mmol) was dissolved in CH₂Cl₂ (28 mL) and triethylamine (0.98 mL) and cooled to 0 °C. MsCl (0.499 mL, 6.45 mmol) was added and the reaction mixture was stirred at 0 °C for 6 h. The reaction mixture was partitioned between water and CH₂Cl₂. The aqueous layer was extracted with
 25

additional CH₂Cl₂ and the combined organic layer was dried with MgSO₄. Evaporation of the solvent afforded 3.0 g of a foam which was purified by sgc (40%-50% EtOAc/hexanes gradient) to give 2.77 g (89%) of Compound IV.


- 5 Compound IV : ¹H NMR (300 MHz, CDCl₃) 8.61-5.97 (m, 2H), 8.40 (d, 8 Hz, 1H), 8.24-8.21 (m, 1H), 7.96 (d, 8 Hz, 2H), 7.86-7.83 (m, 1H), 7.70-7.63 (m, 1H), 7.52 (d, 8 Hz, 2H), 7.46-7.40 (m, 1H), 7.18-7.12 (m, 1H), 4.80-4.70 (m, 1H), 2.71 (s, 3H), 1.56 (d, 7Hz, 3H).



Compound V

- 10 **Compound V.** Compound 7 (26.1 g, 57.4 mmol) was dissolved in CH₂Cl₂ (200mL) and triethylamine (20 mL) and cooled to -78°C. Triflic anhydride (10.45 mL, 62.1 mmol) was added and the reaction mixture was stirred for 3 h. The reaction was quenched with water and the layers were separated. The organic layer was washed with water and brine, then dried with MgSO₄. The solvent was evaporated to give 42 g of a foam. The crude product was purified *via* sgc (33%-50% EtOAc/hexanes gradient) to give 29.7 g (88%) of Compound V.


Compound V : ¹H NMR (300 MHz, CDCl₃) 8.61-8.59 (m, 1H), 8.39 (d, 8 Hz, 1H), 8.29-8.24 (m, 1H), 7.99 (d, 8 Hz, 2H), 7.86-7.82 (m, 1H), 7.67-7.62 (m, 1H), 7.49 (d, 8Hz, 1H), 7.46-7.40 (m, 1H), 7.16-7.10 (m, 1H), 4.89-4.84 (m, 1H), 1.65 (d, 6 Hz, 1H).



Compound VI

- 5 **Compound VI.** Compound V (300 mg, 0.512 mmol) was dissolved in methanol (60 mL). Sodium bicarbonate (720 mg, 8.57 mmol) and 5% palladium on carbon (480 mg) were added. The reaction mixture was shaken on a Parr apparatus under 52 psi of hydrogen gas overnight. The reaction mixture was filtered and the solvent was evaporated. The resulting material was partitioned between EtOAc and aq NaHCO₃.
- 10 The organic layer was dried with MgSO₄ and the solvents were evaporated. The crude product was purified *via* sgc (33% EtOAc/hexanes) to give 257 mg (91%) of Compound VI.


EXAMPLE III



Compound 8


- 15 **Compound 8.** In a flame dried 3-necked flask under N₂ blanket Compound 5 (35.7 g, 91 mmol) was dissolved in anhyd THF (360 mL) and cooled in a dry ice/IPA bath. A solution of n-BuLi (2.5 M in hexanes, 76 mL, 190 mmol) was added at a rate that maintained the internal temperature below -60 °C. The reaction mixture was stirred for 1 h. A solution of 2,6-difluorobenzenesulfonyl fluoride (19.47 g, 99.28 mmol) dissolved in anhyd THF (60 mL) was added. The reaction mixture was stirred
- 20 for 2.5 h, then quenched with saturated aq NH₄Cl (400 mL). EtOAc (500 mL) was added and the layers were separated. The aq layer was extracted with EtOAc and the

combined organic layer was washed with brine and dried with MgSO_4 . The solvent was evaporated to give 60.7 g of an oil which was purified by sgc (15%-40% EtOAc/hexanes gradient) giving 14.4 g (28%) of Compound 8.



Compound 9

Compound 9. Compound 8 (21.1 g, 37.2 mmol) was dissolved in dioxane (47 mL) and aq lithium hydroxide (1.0 M, 41 mL, 41 mmol) was added. After 5.5 h, additional LiOH (20 mL) was added and the reaction mixture was stirred overnight. The reaction mixture was extracted with CH_2Cl_2 , and partitioned between CH_2Cl_2 and water. The aq layer was extracted with additional CH_2Cl_2 and the combined organic layer was dried with MgSO_4 . The solvents were evaporated to give 17.6 g of a foam and the crude product was purified by sgc (1%-3% MeOH (NH_3)/ CH_2Cl_2 gradient) to give 12.2 g (69%) of Compound 9.




Compound VII

Compound VII. Compound 9 (10.7 g, 22.6 mmol) was dissolved in a mixture of CH_2Cl_2 (90 mL) and triethylamine (8 mL) and cooled to -78°C . Triflic anhydride (3.80 mL, 22.6 mmol) was added and the reaction mixture was stirred for 2 h. The reaction was quenched with saturated aq NaHCO_3 and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 . The combined organic layer was

washed with brine and dried with MgSO_4 . The solvents were evaporated and the crude product was purified by sgc to give 9.88 g (73%) of Compound VII.

EXAMPLE IV


Compound 5. In a flame dried flask under N_2 blanket Compound 1 (39.2 g, 132 mmol) was dissolved in anhyd THF (1 L) and cooled in a dry ice/acetone bath. A solution of methyllithium (1.6 M in Et_2O , 82.7 mL, 132 mmol) was added followed by a solution of *n*-BuLi (2.5 M in hexanes, 53 mL, 133 mmol). The reaction mixture was stirred for 25 min and a solution of bis(4-trifluoromethylphenyl) disulfide (46.9 g, 132 mmol) dissolved in THF (200 mL) was added. The reaction mixture was stirred for 2 h then allowed to warm to rt overnight. The reaction was quenched with water and concentrated. The resulting mixture was diluted with EtOAc, washed with water, and dried with Na_2SO_4 . The solvent was evaporated and the crude product was purified via sgc (20% EtOAc/hexanes) to give 49.2 g (95%) of a solid. This material (49.2 g) was dissolved in CH_2Cl_2 (1.2 L) and cooled in an ice bath. MCPBA (60%, 90 g) was added in small portions. After 1 h, the ice bath was removed and the reaction mixture was stirred overnight at rt. The reaction mixture was partitioned between CH_2Cl_2 and 10% aqueous NaHCO_3 . The combined organic layer was washed with water and dried with Na_2SO_4 . The solvent was evaporated and the crude product was purified by sgc (25% EtOAc/hexanes) to give 46.3 g (85%) of Compound 5.



Compound 10


Compound 10. In a flame dried flask under N_2 blanket, Compound 5 (21.55 g, 50.7 mmol) was dissolved in anhyd THF (300 mL) and cooled in a dry ice/IPA bath. A solution of methyllithium (1.6 M in Et_2O , 32 mL, 51 mmol) was added, followed by *n*-BuLi (2.5 M in hexanes, 20.3 mL, 50.7 mmol) and the reaction mixture was stirred for

10 min. A solution of bis-(2-fluorophenyl) disulfide (14.2 g, 55.7 mmol) dissolved in THF was added and the reaction mixture was stirred for 2 h at -78°C . The ice bath was removed and the reaction mixture was allowed to warm to rt and left stirring overnight. The reaction mixture was quenched with saturated aqueous NH_4Cl and extracted with EtOAc. The organic layer was dried with Na_2SO_4 and the solvents were evaporated. The crude product was purified via sgc (25% EtOAc/hexanes) to give 23.2 g of a solid. This material was dissolved in CH_2Cl_2 (400 mL) and cooled in an ice bath. MCPBA (60%, 30.3 g) was added in several portions and the reaction mixture was stirred for 1 h. The ice bath was removed and the reaction mixture was left stirring overnight. The reaction mixture was partitioned between CH_2Cl_2 and 5% aq Na_2CO_3 . The organic layer was washed with water and dried with Na_2SO_4 . The solvents were evaporated and the crude product was purified *via* sgc (25% EtOAc/hexanes) to give 10.84 g (44%) of Compound 10.



Compound 11

Compound 11. Compound 10 (11.88 g, 20.36 mmol) was dissolved in dioxane (200 mL) and aq lithium hydroxide (1.0 M, 400 mL) was added. The reaction mixture was stirred for 3 h then and partitioned between CH_2Cl_2 and water. The organic layer was dried with Na_2SO_4 and concentrated to give 9.34 g (99%) of Compound 11.




Compound VIII

Compound VIII. Compound 11 (0.63 g, 1.29 mmol) was dissolved in a mixture of CH_2Cl_2 (60 mL) and triethylamine (0.27 mL) and cooled in an ice bath. Triflic


anhydride (0.55 g, 1.95 mmol) was added and the reaction mixture was stirred for 1 h. The ice bath was removed, and the reaction mixture was stirred an additional 3 h. The reaction was partitioned between water and CH₂Cl₂. The organic layer was washed with water and dried with Na₂SO₄. The solvent was evaporated and the crude product was purified by sgc (20% EtOAc/hexanes) to give 0.53 g (66%) of Compound VIII.

Compound VIII : ¹H NMR (300 MHz, CDCl₃) 8.89-8.87 (m, 1H), 8.58 (d, 8Hz, 1H), 8.32-8.25 (m, 1H), 8.15-8.11 (m, 1H), 8.03-7.98 (m, 2H), 7.71-7.63 (m, 1H), 7.52-7.48 (m, 2H), 7.47-7.41 (m, 1H), 7.16-7.09 (m, 1H), 5.62 (d, 8 Hz, 1H), 4.90-4.80 (m, 1H), 1.63 (d, 7 Hz, 3H).

EXAMPLE V




Potassium hydroxide (3.1 g, 55.2 mmol), 2-bromo-4-chlorophenol (9.52 g, 45.9 mmol), and 4-fluorobenzonitrile (5.73 g, 47.3 mmol) were added to DMA (25 mL) and the reaction mixture was stirred between 100 °C and 110 °C for one week. The reaction mixture was stirred at rt an additional two days. The solvents were partially removed on the rotary evaporator and the resulting mixture was partitioned between water and a 3:1 Et₂O/hexanes solution. The organic layer was washed with water and brine, then dried with MgSO₄. The solvents were evaporated and the crude product was purified by sgc (20%-30% CH₂Cl₂/hexanes) to give 11.96 g (81%) of an oil.



Compound12


Compound 12. The product of the above step (5.90 g, 19.1 mmol) was placed under N₂ atmosphere and a solution of borane in THF (1.0 M, 21 mL, 21 mmol) was

added causing an exotherm. Once the reaction mixture had returned to rt, it was heated to reflux and stirred at reflux overnight. Additional borane in THF (1.0 M, 20 mL, 20 mmol) was added and the reaction mixture was stirred at reflux for an additional 26 h then allowed to cool to rt. Water (55 mL) was added and the reaction mixture was partially concentrated. The resulting mixture was partitioned between EtOAc and aq NaOH (1.0 M). The organic layer was dried with MgSO₄ and concentrated to give 6.2 g of an oil. This material was dissolved in Et₂O and a solution of HCl in Et₂O was added causing Compound 12 (5.2 g, 78%) to precipitate as a solid.




Compound 13

Compound 13. Compound 12 (5.13 g, 16.6 mmol) was suspended in a mixture of CH₂Cl₂ (40 mL) and triethylamine (7.5 mL). The mixture was cooled in an ice-water bath and TFAA (2.35 mL, 16.6 mmol) was added. The reaction mixture was stirred for 1 h and 20 min and the ice bath was removed. The reaction mixture was stirred for an additional 1 h and 20 min at rt. The reaction mixture was diluted with CH₂Cl₂ (100 mL) and washed with aq citric acid (0.5 M), saturated aq NaHCO₃, water, and brine, then dried with MgSO₄. The solvents were evaporated and the crude product (5.22 g) was purified *via* sgc (10%-20% EtOAc/hexanes gradient) to give Compound 13.




Compound 14

Compound 14. In a flame dried flask under N₂ blanket, Compound 13 (1.00 g, 2.47 mmol) was dissolved in anhyd THF (13 mL) and cooled in a dry ice/IPA bath. Methyllithium (1.4 M in Et₂O, 2.3 mL, 3.22 mmol) was added, followed by n-BuLi (2.5 M in hexanes, 1.3 mL, 3.25 mmol). The reaction mixture was stirred for 1 h at -78 °C. A solution of 2,6-difluorobenzenesulfonyl fluoride (1.10 g, 5.60 mmol) dissolved in THF was added and the reaction mixture was stirred for 4 h. The reaction mixture was quenched with pH 7 sodium phosphate buffer (1.0 M) and EtOAc was added. The layers were separated and the aqueous layer was extracted with additional EtOAc. The combined organic layer was washed with brine and dried with MgSO₄. The solvents were evaporated and the crude product was purified via sgc (20%-33% EtOAc/hexanes) gradient to give 76 mg of Compound 14.



Compound 15


Compound 15. Compound 14 (59 mg, 0.12 mmol) was dissolved in 700 µL of dioxane and LiOH (1.0 M, 300 µL, 0.3 mmol) was added. The reaction mixture was stirred at rt for 24 h then partitioned between CH₂Cl₂ and 1.0 M aq NaOH. The organic layer was dried with MgSO₄ and concentrated. The crude product was purified via PTLC (Merck- silica plates, 3% (MeOH/NH₃)/CH₂Cl₂) to give the desired Compound 15. (21 mg, 45%).




Compound IX

Compound IX. Compound 15 (17 mg, 0.042 mmol) was dissolved in CH_2Cl_2 (166 μL) and DIPEA (20 μL). The flask was cooled in an ice/water bath and MsCl (12 μL , 0.15 mmol) was added. The reaction mixture was stirred at 0 °C for 1 h and 30 min. The resulting mixture was partitioned between water and CH_2Cl_2 . The organic layer was washed with water and brine, then dried with MgSO_4 . The crude product was purified *via* PTLC (50 % EtOAc/hexanes) to give 10 mg (50%) Compound IX.

EXAMPLE VI



Compound 16




Compound X

Compound 16 (0.116 g, 0.22 mmoles) was dissolved in CH_2Cl_2 (4 mL) and cooled to 0 °C. BBr_3 solution (1.0 M in CH_2Cl_2 , 0.66 mL) was added and the ice bath was removed. The reaction mixture was stirred at rt for 48 h and then quenched with water at -78 °C. The reaction mixture was diluted with CH_2Cl_2 and the resulting organic layer was washed with aqueous NaHCO_3 , H_2O (3 X 5 mL), and brine. The organics were dried over Na_2SO_4 and the solvent was removed under vacuum to give 0.09 g of crude product. The product was isolated by PTLC (5% CH_3OH / CH_2Cl_2) to provide Compound X (0.01g, 8.8%).

Compound 16 : ^1H NMR (300 MHz, CDCl_3) 1.54 (d, $J = 6.9\text{Hz}$ 3H), 2.67 (s, 3H), 4.72 (q, $J = 5\text{Hz}$ 1H), 4.86 (br. d, $J = 5\text{Hz}$, 1H, NH), 7.08-8.42 (m, 11H).

5

EXAMPLE VII




10


Compound 17

Compound 17 was converted to Compound 18 using the procedure in example VI.

15




Compound 18



Compound 19

- 5 **Compound 18** (0.34 g, 0.64 mmoles) was dissolved in DMF (11 mL), cesium carbonate (0.84 g, 2.58 mmol) was added and the reaction mixture was cooled to 15 °C. Dry bromodifluoromethane gas was introduced into the solution and bubbled for 15-20 min. Progress of the reaction was monitored by TLC and upon completion the reaction mixture was diluted with EtOAc (20 mL), washed with water (4 X 10 mL), and
10 brine. The organics were dried over Na₂SO₄ and concentrated under reduced pressure to give 0.36 g of an oil. The crude product was purified by PTLC (50% EtOAc/hexanes) to provide 0.31 g (83%) of Compound 19.




Compound XI

Compound 19 was converted to Compound XI using the procedure in example II.


- 20 **Compound XI** : ¹H NMR (400 MHz, CDCl₃) 1.51 (d, J = 7.2 Hz 3H), 2.67 (s, 3H), 4.702 (q, J = 6.8 Hz 1H), 5.05 (br. d, J = 6.4 Hz, 1H, NH), 6.71 (t, J = 11.6 Hz, CF₂H) 7.07-8.47 (m, 11H).

EXAMPLE VIII



5 Compound 20

Compound 20. To a solution of Compound 2 (5.00 g, 12.9 mmol) in anhyd THF (75 mL) at -78°C was added *n*-BuLi (13 mL, 2.5 M in hexanes, 32 mmol) dropwise over 10 min. The reaction mixture was stirred for 30 min. A solution of di-*t*-butyl dicarbonate (3.10 g, 14.2 mmol) in anhyd THF (25 mL) was added in one portion via cannula. The reaction was allowed to proceed for 4 h at -78°C . The reaction mixture was then diluted with EtOAc (~250 mL) and washed successively with saturated aq NaHSO_4 (~100 mL), water (~100 mL), and brine (~100 mL). The organic layer was dried over anhyd MgSO_4 , filtered, and concentrated under reduced pressure to yield a solid. Further purification of the solid by sgc (25% EtOAc/hexanes) gave 5.32 g (84%) of Compound 20 as a solid.




20 Compound XII

Compound XII. Compound 20 (2.06 g, 4.23 mmol) was dissolved in methanol (40 mL) and a solution of potassium carbonate (2.92 g, 21.1 mmol) in water (10 mL) was added. The reaction was allowed to proceed for 18 h. The solvent was then removed by evaporation under reduced pressure. The resulting white solid was partitioned between water (~100 mL) and EtOAc (~400 mL). The aqueous layer was extracted further with EtOAc (~100 mL). The combined organic layers were washed with brine (~500 mL), then dried over anhyd MgSO_4 and filtered. Evaporation of the


solvent gave 1.22 g (74%) of *t*-butyl 2-[(4-(1(*S*)-aminoethyl)phenyl)sulfonyl-5-methoxybenzoate, an oil, which was used in the next step without further purification. MsCl (242 μ L, 357 mg, 3.12 mmol) was added dropwise to a solution of crude *t*-butyl 2-[(4-(1(*S*)-aminoethyl)phenyl)sulfonyl-5-methoxybenzoate (1.22 g, 3.12 mmol) and triethylamine (522 μ L, 379 mg, 3.75 mmol) in anhyd CH_2Cl_2 (3.0 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 5 min, then allowed to warm to rt, and subsequently stirred for 3 h. The reaction mixture was diluted with CH_2Cl_2 (~50 mL) and washed successively with 1 M HCl (~50 mL), water (3 x ~50 mL) and brine (~50 mL). The organic solution was dried over anhyd MgSO_4 , filtered, and concentrated to yield a solid. Subsequent purification of the crude product by sgc (25% EtOAc/hexanes) gave 1.41 g (96%) of Compound XII as a solid.

EXAMPLE IX




Compound 21

Compound 21. In a flame dried flask under N_2 blanket, Compound 5 (400 mg, 1.0 mmol) was dissolved in dry THF (5 mL) and cooled to -78 °C. A solution of *n*-BuLi (1.0 M in hexanes, 1.9 mL, 1.9 mmol) was added and the reaction mixture was stirred for 30 min. 2-Fluorobenzaldehyde (200 mg, 1.6 mmol) was added and the reaction mixture was stirred at -78 °C for 3 h. The reaction mixture was then quenched with saturated aq NH_4Cl (20 mL). Methylene chloride (30 mL) was added and the layers were separated. The organic layer was washed with brine, then dried over Na_2SO_4 , and concentrated to dryness. The crude product was purified via sgc (25% EtOAc/hexanes) to give 330 mg (62%) of Compound 21 as a powder.



Compound 22


- 5 **Compound 22.** Compound 21 (10 mg) was dissolved in CH_2Cl_2 (10 mL). Triethylsilane (40 μL , 0.25 mmol) was added followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (20 μL , 0.16 mmol). The reaction mixture was stirred at rt overnight. After removing the solvent, the crude product was purified via PTLC (25% EtOAc/hexanes) to give 6.0 mg (62%) Compound 22 as an oil.



Compound XIII


- 10 **Compound XIII.** Compound 22 (12 mg) was dissolved in methanol (2 mL) at rt.
- 15 NaOH (1.0 M, 2 mL, 2.0 mmol) was added and the mixture was stirred at rt for 2 h. The solvent was removed, CH_2Cl_2 (15 mL) and brine (15 mL) were added, and the layers were separated. The aqueous layer was extracted with additional CH_2Cl_2 (15 mL) and the combined organic layers were dried over Na_2SO_4 and concentrated to dryness. The crude product was then dissolved in CH_2Cl_2 (10 mL) and cooled to 0 °C.
- 20 MsCl (14 μL , 0.18 mmol) was added followed by addition of pyridine (30 μL , 0.37 mmol). The reaction mixture was slowly warmed to rt and stirred overnight. Brine (15 mL) was added and extracted. The organic layer was dried over Na_2SO_4 and concentrated to dryness. The crude product was purified via PTLC (25% EtOAc/hexanes) to give 10 mg (86%) of Compound XIII as an oil.

EXAMPLE X



Compound 23


Compound 23. Compound 21 (330 mg, 0.64 mmol) was dissolved in CH_2Cl_2 (20 mL) at rt. Celite (450 mg) was added followed by addition of PCC (450 mg, 2.1 mmol). The mixture was stirred at rt overnight. The solid was removed by filtration and the organic layer was washed with aq. NaHCO_3 and brine. The organic layer was dried over Na_2SO_4 and concentrated to dryness. The crude product was purified via sgc (33% EtOAc/hexanes) to give 310 mg (94%) of Compound 23 as a powder.



Compound XIV

Compound XIV. Compound 23 (15 mg) was dissolved in methanol (2 mL) at rt. NaOH (1.0 M, 2 mL, 2.0 mmol) was added and the mixture was stirred at rt for 2 h. The solvent was removed and CH_2Cl_2 (15 mL) and brine (15 mL) were added and the layers separated. The aq layer was extracted with additional CH_2Cl_2 (15 mL) and the combined organic layer was dried over Na_2SO_4 and concentrated to dryness. The crude product was then dissolved in CH_2Cl_2 (10 mL) and cooled to 0°C . MsCl (15 μL , 0.19 mmol) was added followed by addition of pyridine (30 μL , 0.37 mmol). The reaction mixture was slowly warmed to rt and stirred overnight. Brine (15 mL) was added and extracted. The organic layer was dried over Na_2SO_4 and concentrated to


dryness. The crude product was purified via PTLC (33% EtOAc/hexanes) to give 9 mg (62%) of Compound XIV as an oil.



5

Compound 24

Compound 24. Oven dried methyltriphenylphosphonium bromide (430 mg, 1.2 mmol) and LHDMS (1.0 M in hexanes, 1.8 mL, 1.8 mmol) were stirred in dry THF (5 ml) at 0 °C for 20 min., then warmed to rt and stirred for 10 min. A solution of Compound 23 (300 mg, 0.58 mmol) in THF (1 mL) was added dropwise. The mixture was stirred at rt overnight. EtOAc (20 ml) was added and the organic solution was washed with brine. The organic layer was dried over Na₂SO₄ and concentrated to dryness. The crude product was purified via PTLC (25% EtOAc/hexanes) to give 260 mg (87%) of Compound 24 as an oil.



Compound XV


20

Compound XV. Compound 24 (200 mg, 0.39 mmol) was dissolved in methanol (3 mL) at rt. NaOH (1.0 M, 3 mL, 3.0 mmol) was added and the mixture was stirred at 50 °C for 2 h. The solvent was removed, CH₂Cl₂ (20 mL) and brine (20 mL) were added, and the layers were separated. The aqueous layer was extracted with additional CH₂Cl₂ (15 mL) and the combined organic layers were dried over Na₂SO₄ and concentrated to dryness. The crude product was then dissolved in CH₂Cl₂


25

(15 mL) and cooled to 0 °C. MsCl (200 µL, 2.5 mmol) was added followed by addition of pyridine (400 µL, 4.9 mmol). The reaction mixture was slowly warmed to rt and stirred overnight. Brine (15 mL) was added and the organic layer separated, dried over Na₂SO₄ and concentrated to dryness. The crude product was purified via PTLT
 5 (50% EtOAc/hexanes) to give 160 mg (82%) of Compound XV as an oil.

EXAMPLE XI




Compound 25



Compound 26

Compound 26. Compound 25 (1.3 g, 2.7 mmol) was stirred at rt with a mixture of CH₂Cl₂/TFA (2:1, 30 mL) for 3 h. The reaction mixture was then poured into brine (40 mL). The layers were separated. The aq layer was extracted with CH₂Cl₂ (3 X 30 mL) and the combined organic layers were dried over Na₂SO₄ and concentrated to dryness. The crude product was dissolved in CH₂Cl₂ (30 mL). EDCI (0.75 g, 3.9 mmol) and pentafluorophenol (0.73 g, 4.0 mmol) were added and the mixture was stirred at rt overnight. The reaction mixture was extracted with diluted aq NaOH and washed with brine. The organic layer was then dried over Na₂SO₄ and concentrated


to dryness. The crude product was purified via sgc (33% EtOAc/hexanes) to give 1.15 g (72%) of Compound 26 as a foam.



Compound XVI

Compound XVI. Compound 26 (50 mg) was dissolved in CH_2Cl_2 (2 mL). 1-Adamantanamine (21 mg, 0.14 mmol) was added followed by addition of DIPEA (0.05 mL, 0.29 mmol). The reaction mixture was shaken overnight. The reaction mixture was then subjected to Amberlyst 15 resin (300 mg, loading 4.1 mmol/g), and was again shaken overnight. The resin was removed by filtration. The filtrate was subjected to MP carbonate resin (Argonaut Technologies) (100 mg, loading 2.64 mmol/g) for 4 h. The resin was removed by filtration and the filtrate concentrated to give 33 mg (70%) of Compound XVI as a powder.

EXAMPLE XII




Compound 27

Compound 27. Compound 5 (500 mg, 1.3 mmol) was dissolved in dry THF (6 mL) at rt. NaH (53 mg, 60%, 1.3 mmol) was added, and the reaction mixture was stirred at rt for 1 h. The reaction mixture was then cooled to -78°C , and n-BuLi (1.0


M in hexanes, 1.5 mL, 1.5 mmol) was added dropwise under N₂ atmosphere. The reaction was stirred at -78 °C for 40 min. A solution of I₂ (390 mg, 1.5 mmol) in THF (2 mL) was added dropwise. The reaction mixture was stirred at -78 °C for 3 h, then quenched with saturated aq NH₄Cl (20 mL). EtOAc (30 mL) was added and the

5 layers were separated. The organic layer was washed with brine, then dried over Na₂SO₄, and concentrated to dryness. The crude product (640 mg) was used without further purification. The crude product (60 mg) was dissolved in toluene (2 mL) and Pd(OAc)₂ (2 mg), P^tBu₃ (1 drop), NaO^tBu (14 mg, 0.15 mmol) and *p*-Chloroaniline (13 mg, 0.11 mmol) were added. The mixture was kept in a sealed tube and heated to 120 °C for 20 h. After cooling, methylene chloride (30 mL) and brine (20 mL) were added and the layers were separated. The organic layer was washed with brine, then dried over Na₂SO₄, and concentrated to dryness. The crude product was purified with via PTLC (20% EtOAc/hexanes) to give 18 mg (30 %) of Compound 27 as a powder.




Compound XVII

Compound XVII. Compound 27 (12 mg) was dissolved in methanol (2 mL) at
rt. NaOH (1.0 M, 2 mL, 2.0 mmol) was added and the mixture was stirred at rt for 3 h. The solvent was removed, CH₂Cl₂ (20 mL) and brine (20 mL) were added, and the layers were separated. The aq layer was extracted with additional CH₂Cl₂ (15 mL) and the combined organic layers were dried over Na₂SO₄ and concentrated to dryness. The crude product was then dissolved in CH₂Cl₂ (15mL) and cooled to 0 °C.
25 MsCl (15 μL, 0.19 mmol) and pyridine (30 μL, 0.37 mmol) were added. The reaction mixture was slowly warmed up to rt and stirred overnight. Brine (15 mL) was added and the reaction mixture was extracted with CH₂Cl₂. The organic layer was dried over Na₂SO₄ and concentrated to dryness. The crude product was purified via PTLC (33% EtOAc/hexanes) to give 6.0 mg (52%) of Compound XVII as an oil.



Compound 28


- 5 **Compound 28.** Compound 5 (500 mg, 1.3 mmol) was dissolved in dry THF (6 mL) at rt. NaH (53 mg, 60%, 1.3 mmol) was added, and the mixture was stirred at rt for 1 h. The reaction mixture was cooled to -78°C , and *n*-BuLi (1.0 M, 1.5 mL, 1.5 mmol) was added dropwise under N_2 atmosphere, and the temperature was maintained at -78°C for 40 min. A solution of I_2 (390 mg, 1.5 mmol) in THF (2 mL)
- 10 was added dropwise. The reaction was stirred at -78°C for 3 h. The reaction mixture was quenched with saturated aq NH_4Cl (20 mL). EtOAc (30 mL) was added and the layers were separated. The organic layer was washed with brine, then dried over Na_2SO_4 , and concentrated to dryness. The crude product (640 mg) was used without further purification. The crude product (60 mg) was dissolved in toluene (2 mL) and
- 15 NaH (5 mg, 60%, 0.12 mmol), $\text{CuBr}\cdot\text{Me}_2\text{S}$ (34 mg, 0.17 mmol) and *p*-chlorophenol (15 mg, 0.12 mmol) were added. The reaction mixture was kept in a sealed tube and heated to 120°C overnight. After cooling, CH_2Cl_2 (30 mL) and brine (20 mL) were added and the layers were separated. The organic layer was washed with brine, then dried over Na_2SO_4 , and concentrated to dryness. The crude product was purified via
- 20 PTLC (20% EtOAc/hexanes) to give 19 mg (31 %) of Compound 28 as a powder.



Compound XVIII


Compound XVIII. Compound 28 (15 mg, 29 μmol) was dissolved in methanol (2 mL) at rt. NaOH (1.0 M, 2 mL, 2.0 mmol) was added and the mixture was stirred at rt for 2 h. The solvent was removed and CH_2Cl_2 (20 mL) and brine (20 mL) was added and the layers were separated. The aq layer was extracted with additional CH_2Cl_2 (15 mL) and the combined organic layer was dried over Na_2SO_4 and concentrated to dryness. The crude product was then dissolved in CH_2Cl_2 (15 mL) and cooled to 0 $^\circ\text{C}$. MsCl (20 μL , 0.25 mmol) was added followed by addition of pyridine (20 μL , 0.25 mmol). The reaction mixture was slowly warmed up to rt and stirred overnight. Brine (15 mL) was added and extracted with CH_2Cl_2 . The organic layer was dried over Na_2SO_4 and concentrated to dryness. The crude product was purified via PTLC (50% EtOAc/hexanes) to give 7.0 mg (48%) of Compound XVII as an oil.

EXAMPLE XIII




Compound 29

Compound 29. To a solution of N,N,N-Trimethylethylenediamine (1.2 mL, 8.6 mmol) in THF (8 mL) at $-20\text{ }^\circ\text{C}$ was added n-BuLi (1.6 M, 5.4 mL, 8.6 mmol) dropwise. After 15 min 4-trifluoromethoxybenzaldehyde (1.5 g, 7.8 mmol) in THF (8 mL) was added. The mixture was stirred for 15 minutes and additional n-BuLi (1.6M, 14.6 mL, 23 mmol) was added. The reaction mixture was stirred at $-20\text{ }^\circ\text{C}$ for 1h, then placed in the freezer at $-20\text{ }^\circ\text{C}$ for 20 h. The mixture was cooled to $-40\text{ }^\circ\text{C}$, and a solution of bis(2-fluorophenyl)disulfide (4.0 g, 15.7 mmoles) in 30 mL THF was added. The reaction mixture was stirred at $-35\text{ }^\circ\text{C}$ for 3 h. The reaction mixture was poured into 0.5 N HCl and extracted with EtOAc. The organic layer was washed with water and brine, dried over Na_2SO_4 , filtered and concentrated to an oil. Purification by sgc (3 % EtOAc / hexanes) gave 1.55 g (62 %) of Compound 29 as a solid..




Compound 30

- 5 **Compound 30.** Methyllithium (3.25 mL, 5 mmol, 1.4 M ether) was added to a solution of Compound 1 (1.22 g, 4 mmol) at -70°C . After 10 min n-BuLi (1.6 M in hexanes, 2.83 mL, 5 mmol) was added and stirred for 30 min. A solution of Compound 29 (1.44g, 4.55mmoles), dissolved in THF (15mL) was added. The resulting mixture was stirred at -70°C for 2.5 h, quenched with water, warmed to 0°C and then
- 10 extracted with 2 X 50 mL EtOAc. The organic layer was washed with water, dried (Na_2SO_4), filtered and concentrated to an oil. Purification by sgc (EtOAc : hexanes) gave Compound 30 (1.4 g, 58%) as a gum.




Compound 31

- 15 **Compound 31.** Triethylsilane (3.5 mL, 22.5 mmol) was added to a solution of Compound 30 (0.6 g ,1.125 mmol) in CH_2Cl_2 (30 mL), followed by addition of boron
- 20 trifluoride etherate (0.32 mL, 1.94 mmol). After stirring at rt for 15 min the reaction mixture was diluted with 50 mL CH_2Cl_2 , washed with water, dried over Na_2SO_4 , filtered, and concentrated to give a solid. Purification via PTLC (25%EtOAc/hexanes (1:3) gave Compound 31 (0.47 g, 89 %) as a solid.




Compound 32

5 **Compound 32.** MCPBA (1.56 g (56%), 5.09 mmol) was added to a solution of Compound 31 (0.47 g, 0.9 mmol) in CH_2Cl_2 (30 mL) at rt. After stirring for 16 h the reaction was washed with 5% aq NaHSO_3 , aq NaHCO_3 , and water. The organics were dried over Na_2SO_4 , filtered, and concentrated to give Compound 31 (0.4 g, 82%) as a solid.




Compound 33

15 **Compound 33.** 1 M aq LiOH (9.7 mL, 9.7 mmol) was added to a solution of Compound 32 (1.78 g, 3.2 mmol) in 1,4-dioxane (15 mL). The resulting mixture was stirred overnight. The solvent was removed under reduced pressure and the residue was dissolved in 50 mL CH_2Cl_2 and washed with 10 mL brine. The organics were dried over Na_2SO_4 , filtered and concentrated to an oil, which was used in the next step
20 without additional purification.



Compound 34


- 5 **Compound 34.** Triethylamine (0.28 mL, 2 mmol) was added to a solution of Compound 33 (0.18 g, 0.4 mmol) in CH_2Cl_2 at rt, followed by addition of MsCl (0.061 mL, 7.9 mmol) in 0.2 mL CH_2Cl_2 . The mixture was stirred overnight, then washed with 2 X 10 mL water, dried over Na_2SO_4 , filtered, and concentrated to give an oil. The oil was purified via PTLC using EtOAc : hexanes (1:1) as the solvent to give Compound
- 10 34 (0.137g, 65%) as a solid.



Compound XIX


- 15 **Compound XIX.** Triethylamine (0.296 mL, 2.1 mmol) was added to a solution of Compound 33 (0.4 g, 0.9 mmol) in 8 mL of CH_2Cl_2 , cooled to 0°C , followed by addition of a solution of trifluoromethanesulfonic anhydride (0.54 g, 1.9 mmol) in CH_2Cl_2 (5 mL). The mixture was stirred at 0°C for 3 h, washed with water, dried over
- 20 Na_2SO_4 , filtered, concentrated under reduced pressure to give crude Compound XIX. The crude product was purified via PTLC using 33% EtOAc:hexanes to give Compound XIX as a solid (0.32 g, 62%).

EXAMPLE XIV




Compound XX

Compound XX. Triethylamine (0.018 mL, 0.129 mmol) was added to a solution of Compound 33 (0.05 g, 0.11 mmol) in CH_2Cl_2 (1.5 mL) followed by addition of 4-(trifluoromethoxy)benzenesulfonyl chloride (0.02 mL, 0.118 mmol) in CH_2Cl_2 at rt. The stirring was continued for 10 h. The reaction mixture was diluted with 50 mL CH_2Cl_2 , washed with water, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by PTLC (33%EtOAc: hexanes to give Compound XX as a solid (0.048 g, 65%).



Compound XXI


Compound XXI. Triethylamine (0.012 mL, 0.086 mmol) was added to a solution of Compound 33 (0.033 g, 0.073 mmol) in CH_2Cl_2 (1 mL) at -5°C . A solution of acetyl chloride (0.0057 mL, 0.08 mmol) in 0.5 mL CH_2Cl_2 was added. The mixture was stirred overnight at rt. The organics were washed with water, and then dried over Na_2SO_4 , filtered, and then concentrated under reduced pressure. The resulting crude was purified by PTLC (EtOAc) to provide Compound XXI as a solid (0.009 g, 25%).



Compound XXII

Compound XXII. Cyclopentyl isocyanate (0.0135 g, 0.12 mmol) was added as a CH_2Cl_2 solution (0.5 mL) to a solution of Compound 33 (0.05 g, 0.11 mmol) in CH_2Cl_2 (1 mL). The reaction mixture was stirred at rt overnight. The solvent was removed under reduced pressure and the crude product was subjected to PTLC (EtOAc/hexanes 1:2) to provide Compound XXII (0.04 g, 65%).


EXAMPLE XV



Compound XXIII


Compound XXIII. N-Boc-piperazine (0.5 g, 2.68 mmol) was added to a solution of Compound XIX (0.2 g, 0.34 mmol) in CH_3CN (10 mL). The reaction was heated at 80°C for 72 h. Additional N-Boc-piperazine (0.25 g, 1.34 mmol) was added and heated at 80°C for another 16 h. The solvent was removed under reduced pressure and the crude product was purified via PTLC (50% EtOAc:hexanes) to provide Compound XXIII as a solid, (0.096 g, 37%).

EXAMPLE XVI




Compound 35

Compound 35. Pyridinium chlorochromate (0.194 g, 0.899 mmol) was added to a mixture of Compound 30 (0.4 g, 0.75 mmol) and Celite (0.4 g) in CH_2Cl_2 (10 mL) at rt. The mixture was stirred for 18 h, filtered through Celite and concentrated. The crude material was purified via PTLC using 33% EtOAc:hexanes to obtain Compound 35 (0.4 g, 100%).




Compound 36

Compound 36. MCPBA (1.29 g (56%), 4.18 mmol) was added to a solution of Compound 35 (0.4 g, 0.75 mmol) in CH_2Cl_2 (20 mL) and stirred at rt for 18 h. The reaction was washed with 5% aq NaHSO_3 , 5% NaHCO_3 , and water. The organics were dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via PTLC using EtOAc:hexanes (1:1) to provide Compound 36 (0.34 g, 80%).




Compound 37

- 5 **Compound 37.** Compound 36 was converted to Compound 37 using a procedure similar to that described in example II.




10 Compound 38

- 15 **Compound 38.** LHMDS (0.9 mL, 1M solution THF, 0.896 mmol) was added to a suspension of methyltriphenylphosphonium bromide (0.215 g, 0.6 mmol) in anhydrous THF (10 mL) at 0 °C. The mixture was stirred at 0 °C for 20 min, then for 10 minutes at rt. A solution of Compound 36 (0.17 g, 0.3 mmol) in THF (8mL) was added and stirring continued for 10 h at rt. The mixture was diluted with EtOAc and washed with water. The organics were dried over anhydrous Na₂SO₄, filtered and concentrated. The crude product was purified via PTLC using EtOAc:hexanes (1:3) to provide Compound 38 as a solid. (0.09 g, 54%).



Compound XXIV

- 5 **Compound XXIV.** Compound 38 was converted to Compound XXIV using a procedure similar to that described in example II.




10

Compound XXV


- Compound XXV.** Hydroxylamine hydrochloride (0.076 g, 1.09 mmol) was added to a solution of Compound 37 (0.03 g, 0.055 mmol) in pyridine (0.5 mL). The mixture was heated at 80 °C for 24 h. The mixture was cooled to rt and the solvent was removed under reduced pressure. The residue was dissolved in 50 mL CH₂Cl₂ and washed with water and brine. The organics were dried over Na₂SO₄, filtered and concentrated to provide crude Compound XXV, which was purified via PTLC (EtOAc/hexanes, 1:3) to afford Compound XXV as a solid (0.01 g, 33%).

EXAMPLE XVII



Compound 39


Compound 39. In a flame dried flask under N_2 blanket, Compound 2 (4.00 g, 10.32 mmol) was dissolved in anhyd THF (41 mL) and cooled to $-78^\circ C$. A solution of $n-BuLi$ (2.5 M in hexanes, 8.25 mL, 20.6 mmol) was added and the reaction mixture was stirred for 25 min. Bis-4-chlorophenyl disulfide (3.10 g/ 10.8 mmol) was added and the reaction mixture was stirred at $-78^\circ C$ for 3 h then between $-78^\circ C$ and $-10^\circ C$ for 3 h. The reaction mixture was quenched with pH 7.0 sodium phosphate buffer (1.0 M, 50 mL). The reaction mixture was partitioned between EtOAc and water. The organic layer was washed with brine, then dried with Na_2SO_4 and concentrated to dryness. The crude product (5.44 g foam) was dissolved in CH_2Cl_2 (120 mL) and cooled to $0^\circ C$. MCPBA (7.24 g) was added. The ice bath was removed and the reaction mixture was stirred at rt overnight. Aqueous $NaHCO_3$ and CH_2Cl_2 were added and the layers were separated. The organic layer was washed with aq $NaHSO_3$, $NaHCO_3$, H_2O , and brine then dried with $MgSO_4$. The crude product was purified by sgc (35%-40% EtOAc/hexanes gradient) to give 1.86 g (32%) of Compound 39.



Compound 40

Compound 40. Compound 39 (1.52 g, 2.70 mmol) was dissolved in dioxane (9 mL) and cooled to 0°C. LiOH (1.0 M aq, 3 mL, 3 mmol) was added and the reaction mixture was left stirring overnight, during which time it warmed to rt. The solvents were evaporated. CH₂Cl₂ and aq NaOH were added and the layers were separated.


- 5 The aqueous layer was extracted with additional CH₂Cl₂ and the combined organic layer was dried with Na₂SO₄ and concentrated to give 0.85 g (68%) of Compound 40.



10 Compound XXVI

Compound XXVI. Compound 40 (143 mg, 0.307 mmol) was dissolved in dioxane and sulfamide (0.128, 1.33 mmol) was added. The reaction mixture was stirred at reflux for 24 h then allowed to cool to rt and concentrated. The reaction mixture was purified via PTLC (5% MeOH/CH₂Cl₂) giving 54 mg (32%) of Compound XXVI.


Example XVIII



20 Compound 41


Compound 41. In a flame dried flask under N₂ blanket, 1-chloro-4-fluorobenzene (7.36 g, 56.4 mmol) was dissolved in anhyd THF and cooled in a dry ice/acetone bath. n-BuLi (2.5 M in hexanes, 22.5 mL, 56.3 mmol) was added and the

reaction was stirred for 50 min. 2-Fluorobenzene sulfonyl fluoride (10.3 g, 57.8 mmol) was added and the reaction mixture was left stirring overnight, during which time it warmed to rt. Saturated aq NH_4Cl (100 mL) was added, followed by EtOAc (100 mL) and the layers were separated. The organic layer was washed with water and brine, then dried with MgSO_4 . The solvents were evaporated and the crude product was purified via sgc (10% EtOAc/hexanes) to afford Compound 41 (2.55 g, 16%) as a solid.




Compound 42

Compound 42. 4-Mercaptobenzoic acid (0.54 g, 3.50 mmol) was dissolved in DMA (10 mL) and cooled in an ice bath. Sodium hydride (60% suspension in oil, 0.30 g, 7.5 mmol) was added and the reaction mixture was stirred for 20 min. The ice bath was removed and the reaction mixture was stirred for 1 h. The flask was cooled to 0°C again and compound 41 (1.0 g, 3.46 mmol) dissolved in DMA (5 mL) was added. The reaction mixture was stirred at 0 °C for 30 min, then allowed to warm to rt and stirred overnight. The reaction mixture was diluted with CH_2Cl_2 and washed with 5% aq HCl, water, and brine. The organic layer was dried with Na_2SO_4 and the solvents were evaporated. The crude product was purified via sgc (5% MeOH/ CH_2Cl_2) to give Compound 42 as a solid (1.04 g, 71%).




Compound 43

Compound 43. Pentafluorophenol (0.91 g, 4.94 mmol) and Compound 42 (1.04 g, 2.46 mmol) were dissolved in 30 mL of CH₂Cl₂ and EDCI was added. The reaction was stirred overnight and diluted with water and CH₂Cl₂. The layers were separated and the organic layer was washed with water and dried with Na₂SO₄. The crude product was purified via sgc (5% EtOAc/hexanes) to give 0.9 g (62%) of Compound 43 as a solid.



Compound XXVII


Compound XXVII. Compound 43 (0.15 g, 0.25 mmol) was dissolved in CH₂Cl₂ (5 mL). Morpholine (44 mg, 0.51 mmol) and DIPEA (49 mg, 0.38 mmol) were added and the reaction mixture was stirred at rt for 2h. The reaction mixture was diluted with EtOAc and washed with 5% aq NaHCO₃, water and brine. The organic layer was dried with Na₂SO₄ and the solvents were evaporated. The crude product was purified via sgc (50% EtOAc/hexanes) to give 98 mg (77%) of Compound XXVII.



Compound XXVIII


Compound XXVIII. Compound XXXVII (72 mg, 0.146 mmol) was dissolved in CH_2Cl_2 (3 mL) and MCPBA (ca 50%, 0.11 g, ca 0.36 mmol) was added. The reaction mixture was stirred overnight then diluted with CH_2Cl_2 . The reaction mixture was washed with aq Na_2CO_3 and water then dried with Na_2SO_4 . The solvents were evaporated and the crude product was purified via sgc (60% EtOAc/hexanes) to give 61 mg (79%) of Compound XXXVIII as a solid.

Example XIX



Compound 44

Compound 44. Cyclopropyl benzene (48.5 g, 410 mmol), glacial acetic acid (510 mL), and sodium acetate (38.9 g, 474 mmol) were added to a roundbottomed flask. The flask was cooled in an ice-water bath. A solution of bromine (66.3 g, 414 mmol) dissolved in 105 mL of acetic acid was added dropwise over 90 min. The reaction mixture was stirred at temperatures between 0 °C and 10 °C for 5 h. The reaction was then allowed to warm to rt overnight. Hexanes (1300 mL) and water (250 mL) were added. Aqueous NaHSO_3 (1M) was added until the reaction mixture changed from yellow to clear. The layers were separated. The organic layer was washed with water, 1M aq Na_2CO_3 , and brine, then dried with Na_2SO_4 . The solvent was evaporated and the crude product was purified *via* sgc using hexanes as the mobile phase to give 17 g of p-cyclopropylbromobenzene (21%) (Compound 44).




Compound 45

5

Compound 45. A flask was flame dried under N₂ blanket. Compound 44 (10.0 g, 50.7 mmol) was added, followed by dry THF (100 mL). The resulting solution was cooled to -78 °C. A solution of n-butyl lithium in hexanes (2.27 M, 22.35 mL, 50.7 mmol) was added dropwise *via* syringe. The reaction mixture was stirred for 10 min.

- 10 SO₂ gas was bubbled into the reaction mixture until the pH of a reaction mixture sample was <1 when mixed with water. The reaction mixture was stirred for 30 min at -78 °C. The ice bath was removed and the reaction mixture was allowed to warm to rt. The reaction mixture was stirred for an additional 30 min at rt. The reaction mixture was concentrated to afford a solid. CH₂Cl₂ (500 mL) and N-
- 15 chlorosuccinamide (10.2 g, 76 mmol) were added and the reaction mixture was stirred for 4 hrs at rt. Water and CH₂Cl₂ were added and the layers were separated. The organic layer was washed with water and brine, then dried with MgSO₄. The solution was filtered and the solvents were evaporated to give 13.3 g of crude p-cyclopropylbenzenesulfonyl chloride (Compound 45).

20




Compound 46

25


Compound 46. Crude compound 45 (13.3 g) was dissolved in 200 mL of acetone and 60 mL of water. Potassium fluoride (7.12 g, 122 mmol) was added and the reaction mixture was stirred overnight at rt. The reaction mixture was diluted with EtOAc and washed with water. The organic layer was dried with Na₂SO₄, filtered, and

concentrated to dryness to give 9.80 g (97%) of crude p-cyclopropyl benzenesulfonyl fluoride (Compound 46).



Compound 47


Compound 47. A flask was flame dried under N₂ blanket. Compound 1 (44.29 g, 150 mmol) was added, followed by 500 mL of anhydrous THF. The flask was cooled to -78 °C and a solution of n-butyl lithium in hexanes (1.77 M, 154 mL, 272 mmol) was added over 40 min. The reaction mixture was stirred for 1.5 h at -78 °C, then transferred *via* cannula into a solution of crude p-cyclopropylbenzenesulfonyl fluoride (27.2 g, 135 mmol) dissolved in 200 mL of anhydrous THF over 1.5 h. The reaction mixture was stirred for 1h. Water was added, followed by EtOAc. The layers were separated and the organic layer was washed with aq NH₄Cl, water, and brine, then dried with Na₂SO₄. The solvents were evaporated, and the crude product was purified by sgc (25%-33% EtOAc/Hexanes gradient mobile phase) to give 24.5 g (45%) of compound 47.



Compound 48


Compound 48. A flask was flame dried under N₂ blanket. Compound 47 (16.33 g, 41.1 mmol) was dissolved in 400 mL of anhydrous THF and cooled to -78

°C. A solution of n-butyl lithium in hexanes (2.3 M, 35.7 mL, 82.1 mmol) was added dropwise *via* syringe. The reaction mixture was stirred for 1.5 h at -78 °C. A solution of 2, 2'-dithiodipyridine (8.89 g, 41.1 mmol) dissolved in 40 mL of THF was added and the reaction mixture was stirred for 2 h. The cold bath was removed, and the reaction mixture was allowed to warm to rt overnight. The reaction mixture was cooled with an ice-water bath and the reaction was quenched with 10 mL of water. The reaction mixture was diluted with EtOAc and washed with saturated aq NH₄Cl, water, and brine. The organic layer was dried with Na₂SO₄ and concentrated. The crude product was purified *via* sgc using 1:2 EtOAc/Hexanes as the mobile phase giving 15.49 g (74%) of Compound 48.




Compound 49

Compound 49. Compound 48 (15.49 g, 30.6 mmol) was dissolved in 1 L of CH₂Cl₂ and the flask was placed in a rt water bath. MCPBA (22.0 g, ca 74 mmol) was added in portions and the reaction mixture was left stirring overnight at rt. The reaction mixture was diluted with CH₂Cl₂ and washed with 10% aq NaHCO₃, water, and brine, then dried with Na₂SO₄. The solvent was evaporated and the crude product was purified *via* sgc using a 20%-50% EtOAc/Hexanes gradient as the mobile phase. Compound 49 (9.4 g, 57%) was isolated as a solid.



Compound 50


Compound 50. Compound 49 (10.16 g, 18.87 mmol) was dissolved in 300 mL of p-dioxane and 300 mL of 1.0 M aq LiOH was added. The reaction mixture was stirred at rt for 3 h. The reaction mixture was diluted with CH₂Cl₂. The layers were separated, and the organic layer was washed with water and brine, then dried with Na₂SO₄. The solvents were evaporated to give 9.0 g of crude Compound 50.



Compound XXIX

Compound XXIX. Crude compound 50 (7.74 g, 17.5 mmol) was dissolved in CH₂Cl₂ (250 mL). Diisopropylethylamine (2.71 g, 21 mmol) was added and the flask was cooled to -78 °C. A solution of triflic anhydride (5.97 g, 21.1 mmol) dissolved in CH₂Cl₂ (50mL) was added dropwise over 1h. The reaction mixture was stirred for 2 h at -78 °C. The cold bath was removed, and the reaction mixture was allowed to warm to rt overnight. The reaction mixture was diluted with CH₂Cl₂ and washed with water and brine. The organic layer was dried with Na₂SO₄ and the solvents were evaporated. The crude product was purified *via* sgc using 1:2 EtOAc/Hexanes as the mobile phase to give 8.61 g (85%) of Compound XXIX.


Compound XXIX: ¹H NMR (300 MHz, CDCl₃): δ 8.56-8.52 (m, 1H), 8.32-8.21 (m, 3H), 8.02-7.92 (m, 4H), 5.42 (d, 9 Hz, 1H), 8.02-7.92 (m, 4H), 5.42 (d, 1H, 9 Hz), 4.84-4.78 (m, 1H), 2.16-2.06 (m, 1H), 1.60 (d, 7Hz, 3H), 1.20-1.17 (m, 2H), 0.97-0.89 (m, 1H).



Compound XXX

5 **Compound XXX.** Compound XXX was prepared from compound 47 using the procedures in example II.


Compound XXX: ^1H NMR (300 MHz, CDCl_3): δ 8.33-8.22 (m, 3H), 8.00-7.94 (m, 2H), 7.66-7.58 (m, 1H), 7.53-7.37 (m, 4H), 7.16-7.05 (m, 1H), 5.160 (d, 9 Hz, 1H), 4.88-4.83 (m, 1H), 2.17-2.06 (m, 1H), 1.65 (d, 7 Hz, 3H), 1.28-1.20 (m, 2H), 0.97-0.90 (m, 2H).



Compound XXXI


15 **Compound XXXI.** The potassium salt of compound XXIX (56 mg, 0.09 mmol) was dissolved in CH_2Cl_2 (5 mL) and Na_2HPO_4 (0.13 g, 0.91 mmol), and urea-hydrogen peroxide complex (85 mg, 0.90 mmol) were added. Trifluoroacetic acid was added (47 mg, 0.22 mmol) and the reaction mixture was refluxed for 4 h then left stirring overnight at rt. Additional urea-hydrogen peroxide complex (85 mg, 0.9 mmol) and TFAA (0.56 mmol) were added and the reaction mixture was refluxed for 6h. The reaction mixture was allowed to cool to rt and diluted with CH_2Cl_2 and water. The layers were separated and the organic layer was washed with water, dried with Na_2SO_4 , and concentrated. The crude product was purified via PTLC on silica using EtOAc as the mobile phase to give 34 mg (64%) of compound XXXI.

Compound XXXI: ^1H NMR (300 MHz, CDCl_3): δ 8.38-8.29 (m, 2H), 8.17 (d, 8 Hz, 1H), 8.07-8.02 (m, 1H), 7.91-7.85 (m, 2H), 7.56-7.36 (m, 5H), 6.11 (d, 8 Hz, 1H), 4.84-4.78 (m, 1H), 2.12-2.01 (m, 1H), 1.57 (d, 7Hz, 3H), 1.21-1.12 (m, 2H), 0.92-0.86 (m, 2H).



Compound XXXII


Compound XXXII. Compound V (0.50 g, 0.85 mmol), zinc (II) cyanide (65 mg, 0.55 mmol), zinc dust (11 mg, 0.17 mmol), 1,1'-Bis(diphenylphosphino)ferrocene (21 mg, 0.04 mmol), and tris(dibenzylideneacetone) dipalladium (17 mg, 0.129 mmol) were added to a 25 mL flask. Dimethylacetamide was added and the reaction mixture was placed under N_2 blanket and heated to 110 $^\circ\text{C}$. The reaction mixture was stirred at 110 $^\circ\text{C}$ for 4 h, then partitioned between EtOAc and water. The organic layer was washed with 2M ammonium hydroxide, water, and brine, then dried with MgSO_4 . Evaporation of the solvent afforded 0.49 g of an oil that was purified *via* sgc using a 20%-25% EtOAc/Hexanes gradient mobile phase to afford compound XXXII (0.20 g).



Compound XXXIII


Compound XXXIII. Compound V (0.51 g, 0.87 mmol), tris(dibenzylideneacetone) dipalladium (40 mg, 0.04 mmol), 2-(dicyclohexylphosphino)-

biphenyl (36 mg, 0.103 mmol), and sodium tert-butoxide (204 mg, 2.12 mmol) were added to a Schlenck flask under N₂ blanket. Toluene (2.5 mL) was added, followed by benzophenone imine (210 mg, 1.15 mmol). The reaction mixture was stirred overnight at 70 °C under N₂. The reaction mixture was allowed to cool to rt and 1 M aq HCl was added. The reaction mixture was diluted with EtOAc and the layers were separated. The organic layer was washed with water and brine, then dried with MgSO₄. The resulting material was filtered and concentrated to give 0.37 g of an oil. The crude product was purified *via* sgc using a 25%-50% EtOAc/Hexanes gradient mobile phase, followed by a 5%MeOH/45%EtOAc/50%Hexanes mobile phase to give 0.11 g of an oil as product.



Compound XXXIV


Compound XXXIV. Compound V (264 mg, 0.45 mmol), sodium tert-butoxide (103 mg, 1.07 mmol), tris(dibenzylideneacetone) dipalladium (107 mg, 0.116 mmol), and 2-(di-tert-butyl-phosphino)biphenyl (61 mg, 0.20 mmol) were added to a Schlenck flask under N₂. THF (1.5 mL) and cyclopropylamine (0.6 g, 10.5 mmol) were added and the reaction mixture was stirred for 24 h at rt. EtOAc and 1 M aq HCl were added and the layers were separated. The organic layer was washed with 1 M aq HCl, water, and brine, then dried with MgSO₄. Filtration and evaporation of the solvents gave an oil which was purified *via* sgc using 25% EtOAc/Hexanes as the mobile phase. Compound XXIV (109 mg) was obtained as a foam.



Compound XXXV

5 **Compound XXXV.** Compound XXXV was prepared from compound 5 according to the procedures in Example XIX.

10 Compound XXXV: ^1H NMR (300 MHz, CDCl_3): δ 8.88 (d, 1.2 Hz, 1H), 8.51-8.56 (m, 2H), 8.31 (dd, 8 Hz, 1 Hz, 1H), 8.18 (dd, 8 Hz, 1 Hz, 1H), 8.08-7.96 (m, 3H), 7.62-7.48 (m, 3H), 5.51 (d, 9Hz, 1H), 4.90-4.70 (m, 1H), 1.62 (d, 7 Hz, 3H).




15 **Compound XXXVI**

Compound XXXVI. Compound XXXVI was prepared from compound XXXV according to the procedure in Example XIX.


20 Compound XXXVI: ^1H NMR (300 MHz, CDCl_3): δ 10.19 (d, 7.8 Hz, 1H), 8.27-8.42 (m, 4H), 8.13 (dd, 7.8 Hz, 2.1 Hz, 1H), 7.93 (d, 8.4 Hz, 2H), 7.78-7.63 (m, 2H), 7.59 (d, 8.4 Hz, 2H), 4.80 (m, 1H), 1.44 (d, 6.9 Hz, 3H).

25




Compound XXXVII

Compound XXXVII. Compound XXXV (0.312 g, 0.548 mmol) was dissolved in 2 propanol (20 mL) and 1.0 M aq NaOH was added (10 mL). The reaction mixture was stirred at temperatures between 80 °C to 84 °C for six days. The reaction mixture was allowed to cool to rt and partially concentrated. EtOAc was added and the layers were separated. The aqueous layer was acidified with 1 M aq H₂SO₄ and extracted with EtOAc. The combined organic layer was dried with MgSO₄ and concentrated to give 0.29 g of an oil. The crude product was purified via sgc using a 25%-33% EtOAc/Hexanes gradient as the mobile phase. The fraction containing Compound XXVII was repurified via sgc using 3% MeOH/CH₂Cl₂ as the mobile phase to give 0.05 g (15%) of Compound XXXVII as a solid.




Compound XXXVIII

Compound XXXVIII. Compound XXXVIII was prepared from compound XXXV according to the procedure used to prepare compound XXXII.




Compound XXXIX

Compound XXXIX. Compound XXXII (0.10 g, 0.17 mmol) was dissolved in acetone (1.5 mL) and water (1 mL). Potassium carbonate (3 mg, 0.022 mmol) and urea-hydrogen peroxide complex (0.16 g, 1.70 mmol) were added and the reaction mixture was stirred overnight at rt. The reaction mixture was diluted with EtOAc and washed with water. The solvents were evaporated and the crude product was purified via PTLC on SiO₂ using 50% EtOAc/Hexanes as the mobile phase to afford Compound XXXIX (75 mg, 73%) as a solid.




Compound XXXX

Compound XXXX. Compound XXXX was prepared from compound 2 according to the procedures in Example II.




Compound XXXXI

Compound XXXXI. Compound XXXXI was prepared from compound XXXX according to the procedure used to convert compound 16 to compound X.



Compound XXXXII


Compound XXXXII. Compound XXXXI (0.15 g, 0.264 mmol) was dissolved in DMA (5 mL). Potassium iodide (0.22 g, 1.30 mmol), cesium carbonate (0.19 g, 0.58 mmol), and 2-bromopropane (49 mg, 0.398 mmol) were added and the reaction mixture was left stirring at rt over the weekend. EtOAc was added and the reaction mixture was washed with satd. aq NH_4Cl and water. The organic layer was dried with Na_2SO_4 and concentrated. The crude product was purified *via* sgc using 3% $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ as the mobile phase to give 83 mg (51%) of Compound XXXXII.



Compound XXXXIII


Compound XXXXIII. Compound XXXXI (0.10 g, 0.176 mmol) was dissolved in DMF (2 mL). Sodium hydride (7 mg, *ca* 1.2 eq) and bromomethylcyclopropane (26 mg, 0.19 mmol) were added and the reaction was stirred at 50 °C for 4 hr then allowed to cool to rt. EtOAc and water were added, and the layers were separated. The organic layer was washed with water and dried with Na_2SO_4 . The solvent was

evaporated and the crude product was purified via sgc using 33% EtOAc/Hexanes as the mobile phase to give 15 mg (14%) of Compound XXXXIII.




Compound XXXXIV

Compound XXXXIV. Compound XXXXIV was prepared according to the procedure used for Compound XXXXIII using ethyl iodide as the electrophile and stirring the reaction at rt overnight before workup.




Compound XXXXV

Compound XXXXV. Compound XXXXI (0.40 g, 0.70 mmol) was dissolved in DMF (8 mL) and NaH (62 mg, ca 2.2 eq) was added. The reaction mixture was stirred for 30 min. Sodium iodide (0.52 g, 3.46 mmol) and 2-chloroethyl methyl ether (80 mg, 0.85 mmol) were added. The reaction mixture was stirred for 1 h at rt then 5 h at 110 °C. The reaction mixture was allowed to cool to rt. EtOAc and satd aq NH₄Cl were added and the layers were separated. The organic layer was washed with water and dried with Na₂SO₄. Evaporation of the solvent, followed by sgc using 50% EtOAc/Hexanes as the mobile phase, afforded 0.21 g (48%) of Compound XXXXV.




Compound XXXXVI

Compound XXXXVI. Compound 50 (50 mg, 0.11 mmol) was dissolved in CH_2Cl_2 (3 mL) and acetic acid (7 mg). Acetone (6 mg, 0.13 mmol), and $\text{NaBH}(\text{OAc})_3$ (36 mg, 0.169) were added, and the reaction mixture was left stirring at rt overnight. EtOAc was added and the reaction mixture was washed with 10% Na_2CO_3 and water. The solvents were evaporated and the crude product was purified via PTLC on SiO_2 using EtOAc as the mobile phase. The resulting product was dissolved in EtOAc and HCl in Et_2O was added causing a white precipitate to form. The solvent was removed and the precipitate was washed with Et_2O and dried *in vacuo* to give 32 mg (49%) of compound XXXXVI as a solid.



Compound XXXXVII


Compound XXXXVII. Compound XXXXVII was prepared according to the procedure used for compound XXXXVI using cyclopentanone as the carbonyl source.



Compound XXXXVIII

5

Compound XXXXVIII. Compound XXXXVIII was prepared according to the procedure used for compound XXXXVI using cyclohexanone as the carbonyl source.




Compound XXXXIX

10

Compound XXXXIX. Compound XXXXIX was prepared according to the procedure used for compound XXXXVI using cyclopropanecarboxaldehyde as the carbonyl source.


20



Compound XXXXX


25

Compound XXXXX. Compound XXVIII (0.10 g, 0.197 mmol) was dissolved in a solution of borane in THF (1.0 M, 1.0 mL, 1.0 mmol). The reaction mixture was refluxed for 4 h then allowed to cool to rt. The solution was concentrated. Methanol (5 mL) and 1 M aq HCl (5 mL) were added and the resulting solution was stirred for 5 h at rt. The reaction mixture was concentrated and EtOAc was added. The resulting solution was washed with aq NaOH and water, then dried with Na₂SO₄. The solvent was evaporated and the crude product was purified via PTLC using 40% EtOAc/Hexanes as the mobile phase. The product isolated from this step was dissolved in EtOAc, and HCl in Et₂O was added causing a precipitate to form. The solvent was removed and the precipitate was washed with Et₂O and dried *in vacuo* to give 22 mg (21%) of Compound XXXXX as a solid.




Compound 51

Compound 51 was prepared from Compound 2 according to the procedures in Example 11.




Compound XXXXI

Compound XXXXI. Compound XXXXI was prepared from compound 51 according to the procedure used to prepare compound XXXVI.




Compound XXXXXII

Compound XXXXXII. Compound XXXXXII was prepared from compound 51 according to the procedure used to prepare compound XXXXVI using 3-methyl-2-thiophenecarboxaldehyde as the carbonyl source.




Compound XXXXXIII

Compound XXXXXIII. Compound XXXXXIII was prepared from compound 51 according to the procedure used to prepare compound XXXXVI using benzaldehyde as the carbonyl source.




Compound 52

Compound 52. Compound 52 was prepared from compound 2 using the procedures in Example II with benzenesulfonyl fluoride as the initial electrophile.



Compound XXXXXIV


Compound XXXXXIV. Compound 52 (0.29 g, 0.67 mmol), cesium carbonate (0.44 g, 1.35 mmol), tris(dibenzylideneacetone) dipalladium (31 mg, 0.034 mmol), dppp (28 mg, 0.068 mmol), and 2-bromopyridine (0.16 g, 1.01 mmol) were dissolved in 11 mL of toluene under N₂ blanket. The reaction mixture was stirred at 80 °C overnight under N₂, then allowed to cool to rt. CH₂Cl₂ was added and the reaction mixture was washed with 2M aq NaHCO₃, water, and brine. The organic layer was dried with Na₂SO₄ and the solvent was evaporated. The crude product was purified via sgc using EtOAc as the mobile phase. The resulting material was dissolved in EtOAc and a solution of HCl/Et₂O was added. The solvents were evaporated to give 145 mg (42%) of Compound XXXXXIV as a solid.



Compound XXXXXV

Compound XXXXXV. Compound XXXV (0.92 g, 1.67 mmol), was dissolved in methanol (40 mL) and 1.0 M aq NaOH was added (20 mL). The reaction mixture was stirred at 70 °C for 21 h. The reaction mixture was concentrated and extracted with EtOAc. The organic layer was washed with 1 M aq HCl, water, and brine, then dried with MgSO₄. The solvent was evaporated and the crude product was purified via sgc using 25%-33% EtOAc/Hexanes as the mobile phase. Compound XXXXXV (0.82 g, 90%) was isolated as an oil.


Compound XXXXV: ¹H NMR (300 MHz, CDCl₃): δ 8.56 (d, 3.9 Hz, 1H), 8.31-8.22 (m, 2H), 8.124 (d, 2.7 Hz, 1H), 8.05-7.95 (m, 1H), 7.92 (d, 8.4 Hz, 2H), .750-7.45 (m, 1H), 7.92 (d, 8.4 Hz, 2H), 7.27-7.23 (m, 2H), 5.8 (d, NH, 1H), 4.85-4.75 (m, 1H), 3.99 (s, 3H), 1.58 (d, 7.2 Hz, 3H).



Compound XXXXXVI

Compound XXXXXVI. Compound 50 was converted to compound XXXXXVI according to the procedure in Example II.


Compound XXXXXVI: ¹H NMR (300 MHz, CDCl₃): δ 8.56-8.52 (m, 1H), 8.31-8.23 (m, 3H), 8.02-7.90 (M, 4H), 4.87-4.78 (d, 7 Hz, 1H), 4.69 (m, 1 H), 2.66 (s, 3H), 2.16-2.06 (m, 1H), 1.51 (d, 7 Hz, 3H), 1.27 -1.17 (m, 2H), 0.96-0.90 (m, 2H).



Compound 53


Compound 53. 2-fluoro-4-chloroaniline (22.90 g, 151 mmol) was dissolved in 120 mL of AcOH and 80 mL of concentrated HCl was added with stirring. The

reaction mixture was cooled to 0 °C and a solution of NaNO₂ (27.2 g, 0.4 mol) dissolved in 40 mL of H₂O was added over 10 min. The reaction mixture was stirred for 30 min at 0 °C. In a separate flask, 500 mg of CuCl was dissolved in 200 mL of AcOH. The flask was cooled to 0 °C and SO₂ gas was bubbled into the solution for 40 minutes. The contents of the "aniline" flask were added to the contents of the second flask over 20 minutes causing a vigorous evolution of gas. After the addition was complete, the ice bath was removed, and the reaction mixture was allowed to warm to rt. The reaction mixture was poured into 500 g of chipped ice and the resulting solids were collected, washed and dried to give 26.1 g (73%) of compound 53.




Compound 54

Compound 54. Compound 53 (4.0 g, 17.5 mmol) was dissolved in acetone (80 mL) and a solution of potassium fluoride (2.03 g, 35 mmol) in water (40 mL) was added. The reaction mixture was stirred at rt overnight. It was partially concentrated on the rotovap, then partitioned between CH₂Cl₂ and water. Evaporation of the solvent afforded Compound 54 (2.60 g, 70%) as an oil.




Compound 55

Compound 55. Compound 55 was prepared from α-methyl benzylamine using a procedure similar to that used to prepare compound 1. N-Iodosuccinamide was substituted for DBDMH and the product was recrystallized from isopropanol/water.




Compound 56

Compound 56. Compound 55 (4.33g, 12.5 mmol) was dissolved in THF (50 mL) and TMEDA (5.6 mL, 37 mmol) was added. The flask was placed under N₂ blanket and cooled to 0 °C. A solution of isopropyl magnesium chloride (2.0 M in THF, 15 mL, 30 mmol) was added via syringe over 6 min. The reaction mixture was stirred at 0 °C for 1 h. The resulting solution was transferred via cannula into a flask containing compound 53 (15 mmol) in an ice-water bath over 15 min. The reaction mixture was left stirring at 0 °C for 1.5 h. Aq NH₄Cl was added and the reaction mixture was extracted with EtOAc. The combined organic layer was washed with brine and dried with MgSO₄. The solvents were evaporated and the crude product was purified via sgc using 1:4 EtOAc/Hexanes as the mobile phase. Solid compound 56 (3.5 g, 68%) was obtained.




Compound 57

Compound 57. Compound 56 was converted to compound 57 using hydrolysis and sulfonylation procedures similar to those described in Example II.



Compound 58

Compound 58. Compound 57 (0.10g, 0.22 mmol) was dissolved in 1 mL of dioxane and 2-mercaptoimidazole was added (28 mg, 0.28 mmol). Sodium hydride (60% dispersion in mineral oil, 18 mg) was added and the reaction mixture was stirred at 100 °C for 8 h. The reaction mixture was quenched with ice and extracted with EtOAc. The organic layer was dried with MgSO₄ and the solvents were evaporated. The crude product was purified via sgc using a 5:95 MeOH/CH₂Cl₂ mobile phase to give 18 mg (15%) of compound 58 as product.



Compound XXXXXVII

Compound XXXXXVII. Compound 57 was oxidized to compound XXXXXVII using a procedure similar to that used to oxidize Compound XIX to compound XXI.

It will be understood that various modifications may be made to the embodiments and examples disclosed herein. Therefore, the above description should not be construed as limiting, but merely as exemplifications of preferred embodiments. Those skilled in the art will envision various modifications within the pe and spirit of the claims appended hereto.